

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 February 2001 (08.02.2001)


PCT

(10) International Publication Number  
**WO 01/09372 A1**

- (51) International Patent Classification<sup>7</sup>: C12Q 1/42, (74) Agents: KLAWITTER Andrew; Bayer Corporation, 63 North Street, Medfield, MA 02052 et al. (US).  
C07D 219/06
- (21) International Application Number: PCT/US00/20429 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 27 July 2000 (27.07.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/146,648 30 July 1999 (30.07.1999) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): BAYER CORPORATION [US/US]; 100 Bayer Road, Pittsburgh, PA 15205-9741 (US).
- (72) Inventors; and (75) Inventors/Applicants (*for US only*): JIANG, Qingping [US/US]; 14 Stratton Way, Northborough, MA 01532 (US). NATRAJAN, Anand [US/US]; 77 Madeline Road, Manchester, NH 03104 (US). SHARPE, David, J. [US/US]; 61 Cross Street, Foxborough, MA 02035 (US). WONG, Wen-Jee [US/US]; 43 Bemis Street, Weston, MA 02493 (US). LAW, Say-Jong [US/US]; 15 Forbes Road, Westwood, MA 02090 (US).
- Published:**  
— With international search report.  
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: CHEMILUMINESCENT SUBSTRATES OF HYDROLYTIC ENZYMES SUCH AS PHOSPHATASES

WO 01/09372 A1



(57) Abstract: Chemiluminescent substrates of hydrolytic enzymes are disclosed having general Formula Lumi-M-P, where Lumi is a chemiluminescent moiety capable of producing light (a) by itself, (b) with MP attached and (c) with M attached, wherein the different properties of Lumi-M-P and Lumi-M allow them to be distinguished. Lumi includes, but is not limited to, acridinium compounds (e.g. acridinium esters, carboxyamides, thioesters and oxime esters), reduced forms thereof (e.g. acridans), and spiroacridan compounds. M is selected from oxygen, nitrogen and sulfur. P is a group that can be readily removed by hydrolytic enzymes to give Lumi-M and P. The hydrolytic enzyme can be phosphatase, glycosidase, peptidase, protease, esterase, sulfatase and guanidinobenzoatase.

# CHEMILUMINESCENT SUBSTRATES OF HYDROLYTIC ENZYMES SUCH AS PHOSPHATASES

5

## CROSS REFERENCE TO RELATED APPLICATIONS

This application is based on Provisional Application  
10 No. 60/146,648, filed July 30, 1999, which is hereby  
incorporated herein by reference.

## STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

15

N/A

## FIELD OF THE INVENTION

This invention relates to novel chemiluminescent  
compounds that are substrates of hydrolytic enzymes, the  
20 chemiluminescent products of which have distinctly  
different light emission characteristics (i.e. emission  
wavelength, kinetics, or quantum yield). This invention  
also relates to a light-releasing reagent composition that  
reacts preferentially with the chemiluminescent substrate  
25 or with the chemiluminescent product in the mixture of the  
two, to generate a discernible signal that can be  
quantified. This invention further relates to detection  
methods comprising a novel acridinium-based  
chemiluminescent substrate, a hydrolytic enzyme and a  
30 signal-releasing reagent. This invention furthermore

relates to detection devices in conjunction with the use of novel chemiluminescent substrates and hydrolytic enzymes, which include a red-sensitive photomultiplier tube or a charge-coupled device, and a light-filtering device to maximize the detection of chemiluminescent product signal. Further, this invention relates to the use of these novel chemiluminescent substrates in assays to detect, quantitatively or qualitatively, a hydrolytic enzyme of interest that is present either as a label or as a marker of a biological sample. Finally, this invention relates to the process and intermediates for the preparation of these novel chemiluminescent substrates.

#### BACKGROUND OF THE INVENTION

The detection of hydrolytic enzymes has been extensively used in diagnostic assays ranging from immunoassays, nucleic acid assays, receptor assays, and other assays, primarily due to their high sensitivity and non-radioactivity. The hydrolytic enzymes include phosphatases, glycosidases, peptidases, proteases, and esterases. By far, most commonly used are phosphatases and glycosidases. For instance, alkaline phosphatase has been extensively used as a label in various enzyme-linked immunosorbent assays (ELISAs) due to its high turn-over rate, excellent thermal stability and ease of use. Many glycosidases, such as  $\beta$ -galactosidase and  $\beta$ -glucuronidase, have also been used in ELISA due to their very high selectivity for the hydrolysis of their preferred substrates. On the other hand, some hydrolytic enzymes play important functions by themselves in biological

-3-

processes of the human body and microorganisms. Therefore, direct detection of these markers is another important aspect of diagnostics.

In connection with the detection of hydrolytic enzymes, there are three types of substrates: chromogenic, fluorogenic and chemiluminescent substrates. Among them, chemiluminescent substrates offer the best enzyme detection sensitivity due to the intrinsic advantages of higher detectability of chemiluminescent product, or lower substrate and instrumental backgrounds, and less interference from biological samples. Therefore, there has been a steady trend towards developing chemiluminescent substrates and applying them in a variety of diagnostics.

15

#### **Stable dioxetanes**

One class of widely used chemiluminescent substrates for hydrolytic enzymes are stable dioxetanes (Bronstein et al, US 4,931,223, 5,112,960, 5,145,772, 5,326,882; Schaap et al, US 5,892,064, 4,959,182, 5,004,565,; Akhavan-Tafti et al, US 5,721,370). Here, the thermally stable protective group on the phenolic moiety of the dioxetane substrates is cleaved by a hydrolytic enzyme of interest, such as alkaline phosphatase (AP) or  $\beta$ -galactosidase, depending on whether the protective group is a phosphoryl or  $\beta$ -D-galactopyranosidyl group. The newly generated dioxetane phenoxide anion undergoes auto-decomposition to a methoxycarbonylphenoxide in an electronically excited state. The latter then emits light at  $\lambda_{\text{max}} \sim 470 \text{ nm}$ .

25

In an aqueous environment where virtually all biological assays are performed, the decomposition of the dioxetanes produces chemiluminescence in a very low quantum yield, typically about 0.01%, and a slow kinetics with  $t_{1/2}$  1  
5 ~ 10 minutes. This is quite different from the decomposition of the dioxetanes in an organic environment. For instance, the dioxetane having the phenol moiety protected by an acetyl group or a silyl group, upon treatment with a base or fluoride, exhibits quantum yield up  
10 to 25% in DMSO and 9.4% in acetonitrile, respectively, and  $t_{1/2}$  is about 5 sec at 25°C. (Schaap, et al: Tetrahedron Letters, 28 (11), 1155, 1987, and WO 90/07511 A).

Voyta et al (US 5,145,772) disclosed a method of intermolecular enhancement of quantum yield of the  
15 dioxetane products using polymeric ammonium salts, which provide a hydrophobic environment for the phenoxide produced by the enzyme.

Akhavan-Tafti et al reported methods of intermolecular enhancement of quantum yield of the  
20 dioxetane products using polymeric phosphonium salts (US 5,393,469) and dicationic surfactants (US 5,451,347, 5,484,556).

Schaap et al (US 4,959,182, 5,004,565) disclosed another method for increasing quantum yield of the  
25 dioxetane products using fluorescent co-surfactants as energy acceptors. The resonance energy embodied in the excited phenoxide produced by the enzyme is effectively transferred to the fluorescent co-surfactants. Instead of emitting light at  $\lambda_{max}$  470 nm characteristic of the  
30 dioxetane, this system emits light at  $\lambda_{max}$  530 nm as a

result of energy transfer to the highly efficient fluorophore, fluorescein.

Another approach for improving quantum yield of the dioxetanes, disclosed by Schaap in US 5,013,827, is to covalently attach a fluorophore having high quantum yield to the light emitting phenoxide moiety. The resonance energy from the excited phenoxide is intramolecularly transferred to the attached fluorophore. The latter in turn emits light at its own wavelength. It is claimed that such dioxetane-fluorophore conjugates exhibit quantum yield as high as 2%.

Wang et al in WO 94/10258 unveiled a class of electron-rich, aryl-substituted dioxetane compounds in which the aryl group is poly-substituted with a suitable electron-donating group so that intense luminescence is observed.

Akhavan-Tafti et al in US 5,721,370 provided a group of stable chemiluminescent dioxetane compounds with improved water solubility and storage stability. The compounds are substituted with two or more hydrophilic groups disposed on the dioxetane structure and an additional fluorine atom or lower alkyl group.

Schaap et al in US 5,892,064 disclosed a class of chemiluminescent dioxetane compounds substituted on the dioxetane ring with two nonspirofused alkyl groups.

Urdea et al, in EP Application 0401001 A2, described another sub-class of dioxetane compounds that can be triggered by sequential treatment with two different activating enzymes to generate light. The system rests on the principle that the dioxetane substrates have two protecting groups that can be removed sequentially by

different processes to produce an excited phenoxide, and the removal of the first protecting group is triggered by the enzyme used as a label in the assay.

## 5 Luminol substrates

Sasamoto et al [Chem. Pharm. Bull., **38**(5), 1323 (1990) and Chem. Pharm. Bull., **39**(2), 411 (1991)] reported that o-aminophthalhydrazide-N-acetyl- $\beta$ -D-glucosaminide (Luminol-NAG) and 4'-(6'-diethylaminobenzofuranyl)-  
10 phthalhydrazide-N-acetyl-  $\beta$  -D-glucosaminide, both being the non-luminescent forms of luminol, are substrates of N-acetyl-  $\beta$  -D-glucosaminidase. Upon the action of the enzyme on these substrates, luminol or luminol derivative is generated, which then can be detected by triggering  
15 with 0.1 % hydrogen peroxide and a peroxidase (POD) or Fe(III)-TCCP complex catalyst to release a chemiluminescent signal.

## Enzyme-modulated protected enhancer and anti-enhancer

20 US Patent 5,306,621 to Kricka disclosed that light intensity of certain peroxidase-catalyzed chemiluminescent reactions can be modulated by AP that acts on a pro-enhancer or a pro-anti-enhancer. For example, the intensity of a chemiluminescent reaction containing  
25 luminol, horseradish peroxidase and hydrogen peroxide can be enhanced by an enhancer (4-iodophenol) liberated by the enzymatic action of AP on a pro-enhancer (4-iodophenol phosphate), thus enabling AP to be assayed. Alternatively, in the same above reaction where additional  
30 enhancer (4-iodophenol) is present, the light intensity


can be decreased by an anti-enhancer (4-nitrophenol) generated by the enzymatic action of AP on a pro-anti-enhancer (4-nitrophenol phosphate). In the latter format, the presence of AP can be assayed by measuring the reduction in the light intensity.

Similar to the above, an assay using enzyme-triggerable protected enhancer for quantitation of hydrolytic enzymes was also unveiled by Akhanvan-Tafti in EP Application 0516948 A1.

Akhanvan-Tafti et al in WO/9607911 disclosed another method of detecting hydrolytic enzymes based on the principle of the protected enhancer, where the light emitting species is generated from the oxidation of acridan by peroxidase and peroxide.

#### Acridene enol phosphate

Akhanvan-Tafti et al (US 5772926; WO 97/26245) disclosed a class of heterocyclic, enol phosphate compounds represented by the non-luminescent acridene enol phosphate. Upon the reaction with a phosphatase enzyme, it is converted to the dephosphoryl enolate, which reacts with molecular oxygen to produce light (see scheme below). It was also disclosed that the light output was greatly enhanced by the addition of a cationic aromatic compound to the assay system.





**Other chemiluminescent substrates for hydrolytic enzymes**

Vijay in US 5,589,328 disclosed chemiluminescent assays that detect or quantify hydrolytic enzymes, such as alkaline phosphatase, that catalyze the hydrolysis of  
5 indoxyl esters. The assay includes the steps of reacting a test sample with an indoxyl ester and then immediately or within a short time (typically less than about 15 minutes) measuring the resulting chemiluminescence. The resulting chemiluminescence may be amplified by adding a  
10 chemiluminescent enhancing reagent.

Among the above chemiluminescent hydrolytic enzyme methods, the dioxetane system appears to be the most sensitive detection system and therefore has been increasingly used in various assays. Despite its  
15 widespread use, this system has an inherent drawback in that background chemiluminescence in the absence of enzyme is observed due to slow thermal decomposition and non-enzymatic hydrolysis of the dioxetane. Another intrinsic disadvantage is that the phenoxide, once generated by  
20 enzymatic reaction, is extremely unstable and readily undergoes decomposition to release light. In this sense, the phenoxide, the light emitting species, is never "accumulated" during the enzymatic reaction. Therefore, it is one of the major goals of this invention to provide  
25 new chemiluminescent substrates whose derived chemiluminescent products have distinguishable emission profiles and whose total signal can be accumulated during the enzymatic reaction, thereby providing an alternative and sensitive detection method for hydrolytic enzymes.

30

## BRIEF SUMMARY OF THE INVENTION

It is an object of this invention to provide chemiluminescent compounds that are substrates of hydrolytic enzymes. Said chemiluminescent substrates upon  
5 treatment with a hydrolytic enzyme convert to the corresponding chemiluminescent products that have distinctly different light emission characteristics.

It is another object of this invention to provide acridinium-based chemiluminescent compounds that are  
10 substrates of hydrolytic enzymes. Said chemiluminescent substrates contain a phenolic moiety or enol moiety in the molecule that is masked by a group, which is thermally and hydrolytically stable. Upon treatment with a hydrolytic enzyme said chemiluminescent substrates convert to the  
15 corresponding acridinium-based chemiluminescent products that have distinctly different light emission characteristics.

It is also an object of this invention that said chemiluminescent substrates and products have distinctly  
20 different light emission characteristics, thereby allowing the separation or distinction of the signal of the substrate from the signal of the product or vice versa when both substrate and product are present in the same test vessel.

It is another object of this invention that said  
25 chemiluminescent products generated by hydrolytic enzymes have emission maxima different from those of their corresponding chemiluminescent substrates.

It is another object of this invention that said chemiluminescent products generated by hydrolytic enzymes

-10-

have quantum yields different from those of their corresponding chemiluminescent substrates.

It is another object of this invention that said chemiluminescent products generated by hydrolytic enzymes  
5 have light-emitting kinetics different from those of their corresponding chemiluminescent substrates.

It is another object of this invention that said chemiluminescent products generated by hydrolytic enzymes have physical and chemical properties different from those  
10 of their corresponding chemiluminescent substrates. Said physical and chemical properties include, but are not limited to, the fundamental net charge distribution, dipole moment,  $\pi$ -bond orders, free energy, or the apparent hydrophobicity/ hydrophilicity, solubility, affinity and  
15 other properties which are otherwise apparent to those who are skilled in the art.

It is another object of this invention that said chemiluminescent substrates are structurally manipulated such that the distinction of the light emission  
20 characteristics can be further enlarged.

It is yet another object of this invention that chemiluminescent products resulting from the action of hydrolytic enzymes on chemiluminescent substrates do not undergo substantial decomposition during the enzymatic  
25 reaction, and thus can be accumulated until triggered by a light-releasing reagent.

The combined objects and advantages of this invention indicated above are attained by:

-11-

A. Firstly, a chemiluminescent substrate of hydrolytic enzyme having the following general Formula I, as follows:

### Lumi-M-P

5


#### Formula I

where "Lumi" is a chemiluminescent moiety capable of producing light (a) by itself, (b) with MP attached and (c) with M attached. Lumi includes, but is not limited to, chemiluminescent acridinium compounds (e.g. acridinium esters, acridinium carboxyamides, acridinium thioesters and acridinium oxime esters) benzacridinium compounds, quinolinium compounds, isoquinolinium compounds, phenanthridinium compounds, and lucigenin compounds, or the reduced (e.g., acridans) or non-N-alkylated forms (e.g., acridines) of the above, spiroacridan compounds, luminol compounds and isoluminol compounds and the like. M is a multivalent heteroatom having at least one lone pair of electrons selected from oxygen, nitrogen and sulfur, directly attached to the light emitting moiety of Lumi at one end and to P at the other end. (When M alone is attached to Lumi to form Lumi-M, it does, of course, have either a proton or a counterion associated with it or is in the form of an ion.). P is a group that can be readily removed by hydrolytic enzymes, as discussed in more detail hereinafter. The light emitting moiety of Lumi is well known.

-12-

For example, when Lumi is an acridinium compound or luminol, the light emitting moiety is the acridinium nucleus or phthaloyl moiety, respectively.

B. Secondly, an enzymatic reaction having the following general reaction A, as follows:



Reaction A

10 where HE is a hydrolytic enzyme, such as phosphatase, glycosidase, peptidase, protease, esterase, sulfatase and guanidinobenzoatase. Lumi-M-P is a chemiluminescent substrate of a hydrolytic enzyme. Lumi-M is a chemiluminescent product having physical and/or chemical  
15 properties different from those of Lumi-M-P. Said physical and/or chemical properties include emission wavelength, quantum yield, light emission kinetics, fundamental net charge distribution, dipole moment,  $\pi$ -bond orders, free energy, or apparent hydrophobicity/ hydrophilicity,  
20 solubility, affinity and other properties.

C. Light releasing reactions which take place on both Lumi-M-P and Lumi-M.

It is one object of this invention to provide novel light-releasing reagent compositions and reagent addition  
25 protocols for triggering light emission from chemiluminescent substrates and products that surprisingly result in better distinction between the signals of the

-13-

chemiluminescent substrates and products. Said light-releasing reagent compositions can be a single reagent and/or multiple reagents, and addition of said multiple reagents to the reaction vessel can be synchronous or sequential. According to the invention, the light releasing reactions may take place on both Lumi-M-P and Lumi-M, as shown in Formula I.

It is another object of this invention that said light-releasing reagent compositions comprise one or more peroxides or peroxide equivalents, and said peroxides or peroxide equivalents include, but are not limited to, hydrogen peroxide.

It is yet another object of this invention that said light-releasing reagent compositions interact with said chemiluminescent substrate and product differentially so that the differentiation between the two signals is optimized.

It is yet another object of this invention that said light-releasing reagent compositions contain one or more enhancers selected from organic, inorganic or polymeric compounds with a broad range of molecular weights, which differentially enhance the light output from either the substrate or the product.

It is yet another object of this invention that said light-releasing reagent compositions contain also one or more quenchers, blockers or inhibitors selected from organic, inorganic or polymeric compounds with a broad range of molecular weights such that they differentially

-14-

quench, block or reduce the light output from either the substrate or the product.

The combined objects and advantages of this invention indicated above are attained by a light-releasing reagent composition. Said light releasing reagent composition consists of two separate reagents, which are sequentially added to a solution containing the chemiluminescent substrate and/or product. The first reagent contains acidic hydrogen peroxide solution, and the second reagent contains an alkaline solution with one or more detergents. Alternatively, to the advantage of better distinction between the signals of certain chemiluminescent substrates and their products, the first reagent contains an alkaline solution with one or more detergents, and the second reagent contains hydrogen peroxide solution.

It is another object of this invention to provide optimal light detection methods for said chemiluminescent reactions, so that the differentiation between light emissions of the substrate and product is optimized for specific applications. Said optimal detection methods comprise the use of a light detection apparatus, which includes a luminometer, a Charge-Coupled Device (CCD) camera, an X-ray film, and a high speed photographic film. Said luminometer comprises a blue-sensitive photomultiplier tube (PMT), or a red-sensitive PMT, or other PMTs optimized for specific applications. Said optimal light detection methods also include the use of an optical filtering device to block or reduce unwanted light emission either from the substrate or from the product. Said optimal detection methods further include a method

and/or a device for detecting or registering light in a sequential manner that eliminates or reduces unwanted signal from the substrate.

The combined objects and advantages of this invention indicated above are attained by one or more light detection methods. One preferred light detection methods comprises the use of a PMT and a long wave pass filter in an enzymatic reaction to detect the chemiluminescent signal from the product that emits light at a wavelength longer than that of the substrate. Another said light detection method comprises the use of a red-sensitive PMT at low temperature (i.e. below 4°C) and a long wave pass filter to improve the detectability of the chemiluminescent product that emits light at a wavelength longer than that of the substrate. Still, another said light detection method comprises the use of blue-sensitive PMT and a short wave pass filter to detect the decrease of chemiluminescent signal from the substrate whose emission wavelength is shorter than that of the product. Yet, another said light detection method comprises the use of a PMT with or without an optical filter within a fixed light measuring time to detect the decrease of chemiluminescent signal from the substrate whose emission kinetics is faster than that of the product.

It is yet another object of this invention to provide methods and assays comprising the use of one or more said chemiluminescent substrates from Formula I, one or more said hydrolytic enzymes, one or more said light-releasing



-16-

reagent compositions, and one or more said optimal light detection methods.

It is yet another object of this invention to provide methods and assays in using one or more said  
5 chemiluminescent substrates to detect, qualitatively or quantitatively, the presence of one or more said hydrolytic enzymes or enzyme conjugates that are present either as labels or as markers of biological samples.

It is a further object of this invention to provide  
10 method that utilize one or more said chemiluminescent substrates from Formula I, and one or more said labeled hydrolytic enzymes to detect, qualitatively and/or quantitatively, the presence of one or more analytes.

Finally, it is also an object of this invention to  
15 provide synthetic methods and intermediates related to the syntheses of said chemiluminescent substrates.

#### BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWING

Figures 1A-1F show structures of chemiluminescent  
20 substrates (2-phos-acridinium esters), capable of short wavelength emission, and their corresponding chemiluminescent products (2-hydroxyl-acridinium esters), capable of long wavelength emission.

Figures 1G-1L show structures of chemiluminescent  
25 substrates (2-Phos-acridinium esters) capable of short wavelength emission, and their corresponding chemiluminescent products (2-hydroxyl-acridinium esters), capable of long wavelength emission, including spiro compounds (Figures 1K-1L).

Figures 1M-1P show structures of chemiluminescent substrates capable of long wavelength emission and its corresponding products capable of short wavelength emission.

- 5        Figures 1Q-1R show structures of chemiluminescent substrate and its corresponding product, which have different emission kinetics.

Figures 2A-2P are the emission spectra of chemiluminescent substrates (2-phos-acridinium esters) and  
10 their corresponding chemiluminescent products (2-hydroxyl-acridinium esters), as well as spectra of chemiluminescent substrates capable of long emission and their corresponding products capable of short emission, determined by Fast Scanning Spectral System (FSSS). The  
15 emission spectra of chemiluminescent substrate and the corresponding products, which have different light emission kinetics, are also shown.

Figure 3 is a plot of wavelength vs. quantum yield of Hamamatsu photomultiplier tube R268.

- 20        Figure 4 is a plot of wavelength vs. quantum yield of Hamamatsu photomultiplier tube R2228P.

Figure 5 is a plot of wavelength vs. quantum yield of the thinned, back-illuminated Charge Couple Device (thinned CCD).

- 25        Figure 6 is a profile of the transmittance of Corion cut-on long wave pass filter LL650 (Lot No. CFS-002645).

Figure 7 is a profile of the transmittance of Corion long wave pass filter LL700.

-18-

Figures 8 and 9 are plots of stabilities of chemiluminescent product (2-OH-DMAE) in pH 10.5, 100 mM Tris buffer containing 1 mM  $\text{MgCl}_2$  as a function of time.

Figure 10 is a plot of showing the detectability of alkaline phosphatase using 2-Phos-DMAE as the substrate. The enzyme reaction was incubated at 45°C for 0.5 hour. The reaction mixture was flashed with 0.25 N NaOH containing 0.5% CTAB immediately followed by 0.5%  $\text{H}_2\text{O}_2$ . The light output was measured for 2 seconds on MLA-1 equipped with R268 PMT and two LL650 long wave pass filters.

Figure 11 is a plot of showing the detectability of alkaline phosphatase under the same reaction condition as described in Figure 10, with the exception that the light output was measured on MLA-1 equipped with R2228P PMT and two LL650 long wave pass filters, and the unit was pre-cooled to 4°C.

Figure 12 is a plot of showing the detectability of alkaline phosphatase using 2-Phos-DMAE as the substrate. The enzyme reaction was incubated at 45°C for 1 hour. The reaction mixture was flashed with 0.25 N NaOH containing 0.5% CTAB immediately followed by 0.5%  $\text{H}_2\text{O}_2$ . The light output was measured for 2 seconds on MLA-1 equipped with R2228P PMT and a LL700 long wave pass filters, and the unit was pre-cooled to 4°C.

Figure 13 is a plot of showing the detectability of alkaline phosphatase using 4'-Phos-AE (Figure 1Q) as the substrate.

Figure 14 is a plot of the standard curve of TSH immunoassay using 2-phos-DMAE (1) as the chemiluminescent substrate and alkaline phosphatase as the label.

5 DETAILED DESCRIPTION OF THE INVENTION

It has been discovered that certain novel chemiluminescent compounds having a general Formula I are substrates of hydrolytic enzymes, with the structure of Formula I shown as follows:

Lumi-M-P

10

Formula I

Wherein

As shown in Formula I, Lumi is defined as a  
15 chemiluminescent moiety capable of producing light (a) by itself, (b) with MP attached and (c) with only M attached. M is defined as a multivalent heteroatom having at least one lone pair of electrons selected from oxygen, nitrogen and sulfur, wherein M is directly attached to the light emitting  
20 moiety of Lumi at one end and to P at the other end. P is a group that can be readily removed by hydrolytic enzymes. When M alone is attached to Lumi to form Lumi-M, M has either a proton or a counterion associated with it or is in the form of an ion.

25 The chemiluminescent moiety of Lumi includes, but is not limited to, acridinium compounds [including acridinium esters (Law et al, US 4745181), acridinium carboxyamides
















-20-

(Mattingly et al., US 5468646; Kinkel et al, *J. Biolumin. Chemilumin.*, **4**, 136-139, 1989), acridinium thioesters (Kinkel et al, *J. Biolumin. Chemilumin.*, **4**, 136-139, 1989), and acridinium oxime esters (Ghitti et al, PCT WO 98/56765)]

5    benzacridinium        compounds,        quinolinium        compounds,        isoquinolinium compounds, phenanthridinium compounds, and lucigenin compounds, or the reduced (e.g., acridans) or non-N-alkylated (e.g., acridines) forms of the above, spiroacridan compounds (Singh et al, PTC WO 94/02486),

10    luminol compounds, and isoluminol compounds, as shown below.

-21-



-22-

Hydrolytic enzymes that may be used in Reaction A, shown previously, include, but are not limited to,

phosphatases: alkaline phosphatase, acidic phosphatase, phospholipase, phosphodiesterase, pyrophosphatase;

glycosidases:  $\beta$ -galactosidase,  $\alpha$ -galactosidase,  $\alpha$ -(D-)-glucosidase,  $\beta$ -glucosidase,  $\beta$ -glucuronidase,  $\alpha$ -manno-sidase, N-acetyl- $\beta$ -D-glucosaminidase, neuraminidase, cellulase,  $\beta$ -fucosidase;

peptidases and proteases: dipeptidylpeptidases I, II and IV, plasminogen activator, calpain, elastase, trypsin, cathepsins B, C, L and O, urokinase, granzyme A, prostatin, thrombin, trypase, follipain, kallikrein, plasmin, prohormone thiol protease, amyloid A4-generating enzymes, human adenovirus proteinase, kallikrein, HIV protease;

esterases: cholinesterase, lipase; and sulfatase, guanidinobenzoatase.


Being thermally and hydrolytically stable in an aqueous environment, the inventive chemiluminescent substrates undergo readily hydrolytic reaction in the presence of hydrolytic enzymes, and the resulting products, as represented by Lumi-M in Reaction A, are also chemiluminescent. It has also been unexpectedly discovered that the light emission characteristics (emission wavelength, kinetics, and quantum yield) of the products differ significantly from those of their corresponding chemiluminescent substrates due to the

-23-

change in some of the chemical and physical properties of the chemiluminescent products generated by hydrolytic reaction. Those chemical and physical properties include the fundamental net charge distribution, dipole moment,  $\pi$ -bond orders, free energy, or the apparent hydrophobicity/hydrophilicity, solubility, and affinity, etc. As a result, a discernible signal due to one or more of these differences is generated, and thus is useful for qualitative or quantitative determination of the specific hydrolytic enzyme that is involved in the Reaction A, shown previously.

#### 1. Substrates and products having different emission wavelength

One preferred class of chemiluminescent substrates selected from Formula I in this invention is related to novel chemiluminescent acridinium compounds represented in Formula II, shown below, with said compounds capable of being chemically triggered to produce light:



Formula II



As shown in Formula II, P is preferably a group that is thermally and hydrolytically stable in aqueous medium, but is readily removable by a hydrolytic enzyme to form Lumi-M; and M is preferably a multivalent heteroatom  
5 having at least one lone pair electrons selected from oxygen, nitrogen and sulfur, and a strong ability of donating electrons to the acridinium nucleus after the removal of P. Preferably, after the removal of the protective group (P) by a hydrolytic enzyme, M becomes  
10 ionizable in the medium of the reaction to bear a negative charge, thus strongly donating electrons to the acridinium ring system.


As shown in Formula II, R<sub>1</sub> is preferably an alkyl, alkenyl, alkynyl or aralkyl group containing from 0 to 20  
15 heteroatoms; more preferably R<sub>1</sub> is methyl and most preferably R<sub>1</sub> is sulfoalkyl or an alkyl containing one or more hydrophilic groups selected from sulfonate, sulfate, -CO<sub>2</sub>H, phosphonate, ethylene glycol, polyethylene glycol, quaternary ammonium (-N<sup>+</sup>R<sub>3</sub>), or any groups containing one  
20 or more of the above (-N<sup>+</sup>R<sub>3</sub>). The inclusion of the hydrophilic moiety in R<sub>1</sub> serves to increase water solubility of the molecule.

As shown in Formula II, C<sub>1</sub>, C<sub>3</sub>, C<sub>6</sub>, and C<sub>8</sub> positions of the acridinium nucleus may be unsubstituted  
25 or substituted. When one or more of said positions are substituted, the substituents (R<sub>2a</sub>, R<sub>2b</sub>, R<sub>3b</sub>, and R<sub>3d</sub>, respectively) are identical or different, selected from -R, substituted or unsubstituted aryl (ArR or Ar), halides, nitro, sulfonate, sulfate, phosphonate, -CO<sub>2</sub>H, -C(O)OR, cyano (-CN), -SCN, -OR, -SR, -SSR, -C(O)R, -C(O)NHR,  
30

ethylene glycol, or polyethyleneglycol. R is selected from the group consisting of alkyl, alkenyl, alkynyl aryl, and aralkyl containing from 0 to 20 heteroatoms.

As shown in Formula II, the C<sub>4</sub>, C<sub>5</sub>, and C<sub>7</sub> positions of the acridinium nucleus may be unsubstituted or substituted. When one or more of said positions are substituted, the substituents (R<sub>2c</sub>, R<sub>3a</sub>, and R<sub>3c</sub>, respectively) are identical or different and are defined the same as R<sub>2a</sub>, R<sub>2b</sub>, R<sub>3b</sub>, and R<sub>3d</sub>. Alternatively, one of R<sub>2c</sub>, R<sub>3a</sub> and R<sub>3c</sub> may be defined as M-P. In this case, C<sub>2</sub> peri-position may be unsubstituted or substituted. When it is substituted, the substituents may be defined as R<sub>2a</sub>, R<sub>2b</sub>, R<sub>3b</sub>, and R<sub>3d</sub>.


Alternatively, as shown in Formula II, any two adjacent substituents at the acridinium nucleus peri-positions can be linked as in the following examples, so as to form additional unsaturated carbocyclic and/or heterocyclic ring fused to the attached acridinium nucleus, as shown below.



As provided in Formula II, A<sup>-</sup> is a counter ion for the electroneutrality of the quaternary nitrogen of the acridinium compounds, which is introduced as a result of quarternerization of the intermediate acridines with alkylating agents, or due to anionic exchange which occurs during the subsequent synthetic steps or in the following work-up of reaction mixtures and purification of the desired compounds in a liquid phase containing excess amount of other anions. Examples of such counter ions include CH<sub>3</sub>SO<sub>4</sub><sup>-</sup>, FSO<sub>3</sub><sup>-</sup>, CF<sub>3</sub>SO<sub>3</sub><sup>-</sup>, C<sub>4</sub>F<sub>9</sub>SO<sub>3</sub><sup>-</sup>, CH<sub>3</sub>C<sub>6</sub>H<sub>4</sub>SO<sub>3</sub><sup>-</sup>, halide, CF<sub>3</sub>COO<sup>-</sup>, CH<sub>3</sub>COO<sup>-</sup>, and NO<sub>3</sub><sup>-</sup>. A<sup>-</sup> is typically not present if the R<sub>1</sub> substituent contains a strongly ionizable group that can form an anion and pair with the quaternary ammonium cationic moiety.

X is nitrogen, oxygen or sulfur.

When X is oxygen or sulfur, Z is omitted and Y is a substituted or unsubstituted aryl group, and preferably Y is a polysubstituted aryl group of the formula III:



5

Formula III

As shown in Formula III,  $R_4$  and  $R_8$  may be identical or different and identified as an alkyl, alkenyl, alkynyl, alkoxy ( $-OR$ ), alkylthiol ( $-SR$ ), or substituted amino groups that serve to stabilize the  $-COX-$  linkage between the acridinium nucleus and the Y moiety, through steric and/or electronic effect; preferably  $R_4$  and  $R_8$  are short chain alkyl groups having from 1 to 10 carbon atoms, more preferably a methyl group, or at least one of  $R_4$  and  $R_8$  is as defined while the other is a hydrogen or an atom selected from halides.  $R_5$ ,  $R_6$  and  $R_7$  may be identical or different, selected from hydrogen,  $-R$ , substituted or unsubstituted aryl, halides, amino,  $-NHR$ ,  $NR_2$ , quaternary ammonium ( $-N^+R_3$ ), hydroxyl, nitro, nitroso, sulfonate, sulfate, cyano ( $-CN$ ), phosphonate,  $CO_2H$ ,  $-SCN$ ,  $-OR$ ,  $-SR$ ,  $-SSR$ ,  $-C(O)R$ ,  $-C(O)NHR$ ,  $-NHC(O)R$ , ethylene glycol, or polyethyleneglycol. Preferably,  $R_5$ ,  $R_6$  and  $R_7$ , identical or different, are a hydrophilic group selected from

sulfonate, sulfate,  $-CO_2H$ , phosphonate, ethylene glycol, polyethylene glycol, quaternary ammonium ( $-N^+R_3$ ), or any groups containing one or more of the above hydrophilic moiety, which serves to increase water solubility of the molecule.

Any adjacent two groups of  $R_4$  to  $R_8$  in Formula III can form one or more additional fused hydrocarbon aromatic rings or heteroaromatic rings with or without substitutions. The additional fused hydrocarbon aromatic rings and heteroaromatic rings include, but are not limited to, benzene, naphthlene, pyridine, thiophene, furan, and pyrrole, etc.

Alternatively, Formula II can represent another class of Lumi-M-P where Lumi is an acridinium oxime ester (as described in PCT #WO 98/56765 hereby incorporated by reference), and M-P are as defined above. When the Lumi is an acridinium oxime ester, said X, Y, and Z of Formula II are separately defined as follows:

As shown in Formula II, when X is oxygen, Z is omitted and Y is defined above or  $-N=CR_9R_{10}$ , wherein  $R_9$  and  $R_{10}$ , identical or different, and are selected from hydrogen, substituted or non-substituted aryl, alkyl, alkenyl, alkynyl, halide, alkoxyl and aryloxy groups.

As shown in Formula II, when X is nitrogen, then Z is  $-SO_2-Y'$ ,  $Y'$  has the same definition of Y as defined above, and Y and  $Y'$  may be the same or different. Additionally, Y itself can be a branched or straight-chain alkyl containing from 1 to 20 carbons, be halogenated or

unhalogenated, or a substituted aryl, or heterocyclic ring system.

Thus, given the alternatives discussed above, X, Y and Z in Formula II can be defined as follows:


5 X is nitrogen, oxygen or sulfur; such that,

when X is oxygen, Z is omitted and Y is a substituted or unsubstituted aryl group or  $-N=CR_9R_{10}$ , wherein  $R_9$  and  $R_{10}$  may be the same or different and are selected from hydrogen, substituted or non-substituted aryl, alkyl, 10 alkenyl, alkynyl, halide, alkoxyl and aryloxy groups;

when X is sulfur, Z is omitted and Y is a substituted or unsubstituted aryl group;


when X is nitrogen, Z is  $-SO_2-Y'$ ,  $Y'$  being defined the same as Y above; Y is as defined above or can be a 15 branched or straight-chain alkyl containing 0 to 20 carbons, halogenated or unhalogenated, or a substituted aryl, or heterocyclic ring system; and Y and  $Y'$  can be the same or different.

Structurally closely related to chemiluminescent 20 acridinium compounds of Formula II are their reduced forms, chemiluminescent acridan compounds, represented as Formula IV, where all substitutions are as defined in Formula II. These compounds can be chemically triggered to produce light via formation of acridinium intermediates 25 and/or via direct oxidation to form electronically excited products, which are the same products of acridinium compounds of Formula II.



Formula IV

Another class of chemiluminescent substrates structurally related to acridinium substrates of Formula II are spiroacridan compounds having Formula V, wherein  $X_1$  and  $X_2$  may be identical or different, are selected from the group consisting of oxygen, sulfur and nitrogen, and when  $X_1$  and/or  $X_2$  are oxygen or sulfur,  $Z_1$  and/or  $Z_2$  are omitted, when  $X_1$  and/or  $X_2$  are nitrogen,  $Z_1$  and/or  $Z_2$  are hydrogen, alkyl, aryl or  $-SO_2-Y'$ ; G is a group connecting  $X_1$  and  $X_2$  to form a ring having 5 to 10 members; and  $R_1$ ,  $R_{2a-c}$ ,  $R_{3a-d}$ , M and P are defined in Formula II.




Formula V

15

Within the chemiluminescent spiroacridan substrates of Formula V is a subclass shown as Formula VI below, where G is singly or multiply substituted ( $R_{11}$ ) or unsubstituted aromatic ring with 0-3 heteroatoms, wherein

5  $R_{11}$  is the group selected from hydrogen, -R, substituted or unsubstituted aryl (ArR or Ar), halides, nitro, sulfonate, sulfate, phosphonate, -CO<sub>2</sub>H, -C(O)OR, cyano (-CN), -SCN, -OR, -SR, -SSR, -C(O)R, -C(O)NHR, ethylene glycol, or polyethyleneglycol, and said subclass of substrates

10 having Formula VI:




Formula VI

Preferably, novel chemiluminescent acridinium

15 substrates of the present invention of Formula II are a subclass of compounds represented by Formula VII below, wherein multivalent heteroatom M is oxygen (O), and group P is a phosphoryl group, -PO<sub>3</sub>Na<sub>2</sub>, where two sodium cations, whose purpose is solely for the electroneutrality of the

20 molecule, can be exchanged independently with hydrogen, potassium, magnesium, calcium, or other cationic ion(s) or group(s).






Formula VII

5        The preferred hydrolytic enzyme (HE) as shown in Scheme I is a hydrolytic phosphatase. Particularly preferred is the use of hydrolytic phosphatase with a compound defined by Formula VII.

10       Structurally closely related to preferred chemiluminescent acridinium compounds of Formula VII are their reduced forms, chemiluminescent acridans, represented as Formula VIII, where all substitutions are as defined in Formula VII. The compounds of Formula VII may be chemically triggered to produce light via formation of acridinium intermediates and/or via direct oxidation to

15       form electronically excited products, which are the same products of acridinium compounds of Formula VII.


-33-



Formula VIII

5

Another subclass of the compounds structurally related to preferred chemiluminescent acridinium compounds of Formula VII are spiroacridan compounds of Formula IX, wherein,  $X_1$ ,  $X_2$ ,  $Z_1$ ,  $Z_2$ ,  $R'$ ,  $R_1$ ,  $R_{2a-c}$ ,  $R_{3a-d}$ ,  $M$  and  $P$  are as defined in Formula VI, and two sodium cations are as defined in Formula VII.




Formula IX

15

One of the preferred compounds selected from Formula VII, in which  $R_{2a-c}$ ,  $R_{3a-d}$ , are hydrogen,  $R_1$  is methyl,  $X$  is

-34-


oxygen, z is omitted, and Y is a polysubstituted aryl moiety of Formula III where R<sub>4</sub> and R<sub>8</sub> are methyl, R<sub>5</sub> and R<sub>7</sub> are hydrogen, and R<sub>6</sub> is carboxyl (-CO<sub>2</sub>H), is 2-Phos-DMAE, whose structure is shown as 1. (See Fig. 1A. Note:  
5 Structure 1 below is also shown in Fig. 1A, attached. Similarly, other structures in this application are referred to by numbers which correspond to the appropriate sequence number in Figure 1.)



1

As disclosed in the section of Examples, 2-Phos-DMAE (1) is an excellent substrate of hydrolytic alkaline  
15 phosphatase (AP). In alkaline aqueous medium of a wide pH range, it is readily dephosphorylated by AP to form 2-OH-DMAE (2) as illustrated in Reaction B. Both 1 and 2 are chemiluminescent. As disclosed in Figures 2A - 2 B and also in Table 1, it has been unexpectedly discovered that  
20 1 and 2 emit light at different emission maxima when they are treated with hydrogen peroxide in strong alkaline solution, respectively. Specifically, compound 1 emits a strong, visible blue light at  $\lambda_{\text{max}}$  478 nm while compound 2

emits a strong, visible orange light at  $\lambda_{\text{max}}$  602 nm, thus resulting in a bathochromic shift of emission maximum by 128 nm. Several pairs of analogs (3 and 4, 5 and 6, 7 and 8), as well as a pair of spiroacridans (11 and 12), whose structures are given in Figure 1, also have the same light emission characteristics. Compounds 7 and 8 also carry additional hydrophilic group on the nitrogen of the acridinium nucleus.



## Reaction B

15 As illustrated in Reaction C, the chemiluminescent reaction of acridinium compound is triggered by hydrogen peroxide in strong alkaline solution. Formation of the high energy dioxetanone follows the departure of the leaving group LG. The highly strained dioxetanone  
20 intermediate decomposes to the acridone, a portion of which is formed in an electronically excited state. When the excited acridone reverts to a ground state, light


emission occurs. The emission wavelength is determined by the energy gap between the first excited state and the ground state, which in turn is determined by specific structure of the acridone having various functional group

5 T. As disclosed in WO 00/09487, which is fully incorporated herein by reference, there are several factors which can influence the energy gap between the first excited state and the ground state of the acridone, and thus affect the emission wavelength. One of the

10 factors is the direct attachment of the requisite functional group T to one of the peri-positions of the acridinium nucleus. It has been discovered and disclosed in the aforementioned WO 00/09487 that when a hydroxyl group is placed at the (2) or (7) position, such as in 2-

15 OH-DMAE (2), the compound emits light at  $\lambda_{\text{max}}$  604 nm in strong alkaline solution. This is because in a strong alkaline solution, the hydroxyl group becomes deprotonated, and the resulting negatively-charged oxy anion exerts a strong electron donating effect to the

20 acridinium nucleus, consequently decreasing the energy gap between the first excited state and the ground state of the acridone, which causes a bathochromic shift of light emission to 602 nm from 430 nm of the unsubstituted acridinium compound.



Reaction C


5        In this invention, a phosphoryloxy group at the (2)  
 position of **1** serves as a masked hydroxyl group. This  
 phosphoryloxy group is thermally and hydrolytically stable  
 in aqueous medium with a wide range of pHs where virtually  
 all hydrolytic enzymatic reactions take place. It is also  
 10    stable in a mixture of hydrogen peroxide and strong  
 alkaline solution for a length of time that is needed for  
 converting acridinium compounds to the light emitting  
 acridones. Because the phosphoryloxy group in **1** is  
 virtually intact in the absence of hydrolytic enzyme  
 15    during the reaction and also stable to hydrogen peroxide  
 and strong alkaline solution, the electron-donating effect  
 of the directly attached oxygen at the (2) position  
 towards the acridinium nucleus is significantly  
 diminished. Thus, no bathochromic shift of emission  
 20    wavelength occurs. This is in sharp contrast to the  
 situation of 2-OH-DMAE (**2**), the product of the reaction in  
 the presence of hydrolytic enzyme. In the latter case,

the hydroxyl group at the (2) position is ionized to a negatively-charged oxy anion, which has a strong electron donating effect to the acridinium nucleus, thus causing a significant bathochromic shift of emission wavelength.

5           It would be desirable that in a system using 1 as a chemiluminescent substrate, a specific hydrolytic enzyme such as AP that is either used as a label or present as a marker of a biomolecule converts a short wavelength emitting 1 to a long wavelength emitting 2. Indeed, upon  
10 treatment with hydrogen peroxide and strong alkaline solution, the reaction mixture containing both 1 and 2 produces a mixture of light signals each with its own maximum at 478 nm and 602 nm, respectively. The decrease in the short signal ( $\lambda_{\text{max}}$  478 nm) or the increase in the  
15 long signal ( $\lambda_{\text{max}}$  602 nm) directly correlates to the amount of the enzyme.

Another preferred compound selected from Formula VII is 2-Phos-7-MeO-DMAE (9). Unexpectedly, the dephosphorylated product, 2-OH-7-MeO-DMAE (10) in Reaction D was  
20 found to have 60-70 nm longer emission maximum than the unmethoxylated compound 2 (2-OH-DMAE), while both of the corresponding 2-phosphorylated forms have nearly the same light emission maxima. Thus the 2-Phos-7-MeO-DMAE and 2-OH-7-MeO-DMAE pair of chemiluminescent substrate and  
25 product provide effectively further improved distinction in light emission wavelength. The discovery has led to another preferred set of acridinium-based chemiluminescent substrates, wherein the Formula VII is further defined as above, except the substituents of  $R_{3c}$  can be any electron-  
30 donating group such as hydroxy, thiol, amino,

monoalkylamino, dialkylamino, alkoxy or thioalkyl. The presence of this second electron-donating group further decreases the energy gap between the electronically-excited and ground states of the light-emitting acridone. Consequently, a further red-shift in light emission is observed compared to 2-hydroxy-DMAE. This shift also results in increased spectral distinction of the blue and red-emitting acridinium ester pair.



10

9

10

## Reaction D

One of the modes of enzyme detection disclosed in this invention is to detect long wavelength signal of the product (2) that arises from enzymatic action. The commonly used methods for light detection include the use of a luminometer, a CCD camera, X-ray film, and a high speed photographic film. Since the hydrolytic enzyme to be detected often exists in a small quantity in the sample, the amount of the product that is generated by the enzyme is relatively small in comparison with the amount of the substrate used in the reaction. Therefore, it is



critical to be able to detect such a small amount of this signal in the presence of a strong signal of the substrate.

One important aspect of this invention related to  
5 detecting a long wavelength emission signal of the product  
such as 2 produced by a hydrolytic enzyme is the use of a  
luminometer having a red-sensitive photomultiplier tube  
(PMT). (Frequently, this type of PMT is cooled during use  
in order to reduce system noise.) The most commonly used  
10 commercial photomultiplier tubes are made from low dark  
count bialkali material. It has an excellent quantum  
yield in the short wavelength region but a very low  
quantum yield in the long wavelength region. For example,  
bialkali PMT model R268 from Hamamatsu has a quantum yield  
15 of about 22% in the range of 400-500 nm and only ~2% at  
600 nm and above, which is not desirable for this specific  
application. On the other hand, PMT made from multialkali  
material is relatively more red-sensitive than that made  
from bialkali material. For instance, multialkali PMT  
20 model R2228 from Hamamatu has a quantum yield of 3~5% over  
the region of 400 to 500 nm, and a quantum yield of ~5%  
around 600 nm. This is more desirable for this  
application. The drawback of this PMT is a high dark  
count, and therefore it has to operate at low temperatures  
25 in order to suppress the dark count. Figures 3 and 4 show  
quantum efficiencies of PMTs of bialkali R268 and  
multialkali R2228P, respectively.

Another useful aspect of this invention related to  
detecting a long wavelength emission signal of the product  
30 is the use of a charge-coupled device (CCD) camera,

particularly a thinned, back-illuminated cooled CCD. As shown in Figure 5, said CCD has a ~80% quantum efficiency at 400 nm and ~90% at 700 nm, thus capable of increasing the detectability of long wavelength emission (> 550 nm) signal by 10-20 times or more over R268.

Since both the substrates and products of the present invention are chemiluminescent, an essential aspect of this invention related to detecting a long wavelength emission signal of 2 is the use of a filtering device to block the short wavelength signal from 1. Figures 6 and 7 show transmittances of Corion LL700 and LL650 long wave pass filters (Corion, Franklin, MA), respectively.

One of the advantages in using chemiluminescent acridinium substrates like 1 to detect hydrolytic enzymes is that the products like 2 generated by the enzyme can be accumulated without undergoing significant decomposition during the enzymatic reaction. Figures 8 and 9 shows the stability of 2-OH-DMAE (2) in aqueous medium at pHs 9 and 10.5 and at different elevated temperatures.

The sensitivity relating to the detection of a long emission signal of the product generated by a hydrolytic enzyme is largely dependent on the signal differentiation of the product (2) and the substrate (1). It is therefore desirable to reduce background emission of the substrate 2-Phos-DMAE in order to obtain a sensitive system. The generation of light from acridinium compounds is traditionally made by the treatment with acidic hydrogen peroxide solution first, followed by an alkaline surfactant solution. The purpose of the acid is to

-42-

convert the pseudo-form of acridinium compound to the quaternary form and the surfactant aids in enhancing the quantum yield of the acridinium compound. Preferably, said acidic hydrogen peroxide solution contains hydrogen  
5 peroxide at concentration of 0.001 to 5%, and nitric acid at concentration of 0.001 to 1N, and said alkaline surfactant solution contains sodium hydroxide at concentration of 0.01 to 1 N, and AQUARD (CTAB) at 0.01 to 1%. More preferably, said acidic hydrogen peroxide  
10 solution contains 0.5% hydrogen peroxide and 0.1 N nitric acid, and said alkaline surfactant solution contains 0.25 N sodium hydroxide and 0.5% AQUARD (CTAB).

An unexpected finding, to the advantage of detecting the long wavelength emission signal from 2, is that under  
15 certain conditions the chemiluminescence from 1 is selectively and significantly suppressed, and thereby the overall signal differentiation of 2 over 1 is improved. Specifically, the chemiluminescence of 1 is significantly lowered if the solution is treated with an alkaline  
20 solution first and then with hydrogen peroxide solution. More specifically, when a solution of 1 is treated with 0.25 N sodium hydroxide solution containing 0.5% AQUARD, followed by 0.5% hydrogen peroxide, the quantum yield of 1 is lowered by more than 30 folds. In contrast, under the  
25 same condition the quantum yield of 2 is basically not affected. This results in an overall improvement of the signal differentiation by 30 folds. Thus, to the advantage of better distinction between the signals of certain chemiluminescent products and their substrates,  
30 preferably, said alkaline solution contains sodium

hydroxide at concentration of 0.01 to 1 N, and AQUARD at concentration of 0.01 to 1%, and said hydrogen peroxide solution contains hydrogen peroxide at concentration of 0.001 to 5%. More preferably, said alkaline solution  
5 contains 0.25 N sodium hydroxide and 0.5% AQUARD, and said hydrogen peroxide solution contains 0.5% hydrogen peroxide.

With one or more of the above approaches combined, several assay systems based on the principle of detecting  
10 the long emission signal generated by the action of AP on 2-Phos-DMAE (1) are constructed, and disclosed in the section of Examples. One example (Example 17) consists of the incubation of AP standards in a substrate solution containing 0.1 mM 2-Phos-DMAE (1), 1 mM magnesium chloride  
15 in 100 mM, pH 9 Tris buffer at 45°C for 1 hour. The light outputs were measured on a luminometer equipped with a red-sensitive PMT (R2228P) and a long pass filter (LL700), which is placed in a 4°C cold room. The sample was treated first with 0.25 N sodium hydroxide solution containing  
20 0.5% AQUARD, immediately followed by 0.5% hydrogen peroxide. As shown in Figure 12, AP is detected at a level below 1.0E-18 mole. Another example under the similar condition is given in Example 16 and the result is shown in Figures 10 and 11.



25 Another important method for selectively reducing background of 2-Phos-DMAE (1) is through a selective quenching. Selective quenching, herein and hereafter, refers to the use of one or more compounds or chemical moieties to selectively reduce chemiluminescence of the  
30 substrate such as 1 via the mechanism of energy transfer.

Referring back to Scheme III, the energy of the excited acridone (donor) resulting from the chemiluminescent reaction of the acridinium compound can be transferred to an adjacent, non-fluorescent molecule (acceptor or quencher). This non-fluorescent molecule reverts to the ground state via a non-radiative pathway to release the energy. The quantum yield of the acridinium compound is quenched or reduced as the result of the quenching. The effectiveness of the quenching depends on the distance, spectral overlap and transition dipole-dipole interaction of the acridinium compound and the quencher. First, the acridinium compound and quencher must be in a close vicinity, preferably at a distance less than 10 nm in order to achieve 20~100% quenching effect, since the effectiveness of energy transfer is inversely proportional to the sixth power of the distance between the donor and acceptor. Secondly, the UV absorption spectrum of the quencher must overlap, at least partially, with the emission spectrum of the acridinium compound. Lastly, even though it is often not easy to control the spatial conformations of the two concerned molecules in order to obtain the effective transition dipole-dipole interaction, a relatively free movement of two spatially close molecules often fulfil this requirement.

There are in general two ways of quenching, intermolecular and intramolecular. Specifically related to this invention, the intermolecular quenching refers to when the quencher and acridinium compound coexist in the same solution, but the two molecules are not linked together. The intramolecular quenching refers to when

both the quencher and acridinium compound are covalently linked in one molecule. In the case of intermolecular quenching, the effectiveness of the quenching depends on the concentration of the quencher relative to the concentration of acridinium compound. The concentration of the quencher or substrate or both must be in the range of millimolar in order to achieve effective quenching. However, for the intramolecular quenching, the effectiveness of the quenching is determined by a bonding distance between the two molecules, and that in turn can be warranted by the proper selection of the length of the tether that connects the two.

Another aspect closely related to the selection of a quencher is that the quencher should only selectively or preferentially quenches light of 2-Phos-DMAE (1) and have little or no effect on the light emission of 2-OH-DMAE (2). To achieve this, the UV absorption spectrum of the quencher should have a maximum overlap with emission spectrum of 1 but a minimum overlap, or preferably no overlap at all, with the emission spectrum of 2. The other criteria for an effective quencher for this application include a large molar extinction coefficient and adequate water solubility. Examples of compounds that are suitable to serve as quenchers are given below, but are not limited to those listed.




It is an intention of this invention to provide, but not limiting to, one mechanism to selectively reduce the light emission of the substrate via selective quenching. It is understandable that any other approach or mechanism  
5 capable of selectively reducing the quantum yield of the chemiluminescent substrates will also fall in the scope of this invention.

The second mode of light detection involves detecting the short emission signal of the substrate (1) that  
10 diminishes as the result of the action of enzymatic action. This mode of detection takes advantage of the high quantum yield of bialkali, blue-sensitive PMTs as discussed earlier. In connection with the use of a blue-sensitive PMT, a filtering device that is capable of  
15 blocking a long emission signal must be also used in order to block long-wavelength light emission of the product in the hydrolytic enzymatic reaction.

Belonging to Formula II, another important sub-class of chemiluminescent substrates related to this invention  
20 are substrates capable of "reverse emission", where Lumi-M-P is capable of emitting long wavelength light while the product (Lumi-M) is capable of emitting short wavelength light. One major advantage in this reverse emission mode is that the product is detected at a short emission signal  
25 with bialkali, blue-sensitive PMT fitted with a short wave pass filter, where quantum yield of the PMT is as high as 22%.

Chemiluminescent substrates capable of the reverse emission are represented by Formula II where M-P is replaced by Formula X:



Formula X

As provided in Formula X, preferably M and P are defined in Formula II. ED is an electron-donating group, preferably an ionizable group, which donates an electron pair to the conjugated system and is selected from hydroxyl, -OR, -NR'R'', thiol (-SH), -SR, and -CHW<sub>n</sub> where n = 1 or 2, W is an electron withdrawing group including but not limited to nitro, nitroso, cyano (-CN), -CHO, -C(O)R, -N<sup>+</sup>RR'R'', -CO<sub>2</sub>H, -CO<sub>2</sub>R, -S(O)R, -SO<sub>2</sub>R, -SO<sub>2</sub>OR, -SO<sub>2</sub>NHR, -SO<sub>2</sub>NR'R'', -SO<sub>3</sub>H, or F, where R is defined in Formula II, R' and R'' are hydrogen or low alkyl, and R, R' and R'' can all be the same or different. ED is interchangeable with R<sub>12</sub> or R<sub>15</sub>.


R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub> and R<sub>15</sub> may be identical or different, and are selected from hydrogen, -R, hydroxyl, amino, halides, nitro, nitroso, sulfonate, sulfate, phosphonate, -CO<sub>2</sub>H, cyano (-CN), -SCN, -OR, -SR, -SSR, -C(O)R, and -C(O)NHR, and R is defined in Formula II; alternatively, any



-48-


adjacent two groups of  $R_{12}$  to  $R_{15}$  can form one or more additional fused hydrocarbon aromatic rings or heteroaromatic rings with or without substitutions, and the additional fused hydrocarbon aromatic rings and  
 5 heteroaromatic rings include, but are not limited to, benzene, naphthlene, pyridine, thiophene, furan, and pyrrole, and so on.

Preferably,  $R_{12}$ ,  $R_{13}$ ,  $R_{14}$  and  $R_{15}$  are all hydrogen, and ED is hydroxy, and chemiluminescent substrates capable of  
 10 reverse emission have Formula XI:





Formula XI

Alternatively, the group represented by Formula X can  
 15 be located at the C<sub>3</sub> position of acridinium nucleus, having Formula XII below:





Formula XII

More preferably, in Formula XI and XII, M is oxygen; P is  
 5 a phosphoryl group,  $-\text{PO}_3\text{Na}_2$ , where two sodium cations can  
 be exchanged independently with hydrogen, potassium,  
 magnesium, calcium, or other cationic ion(s) or group(s)  
 for maintaining the electroneutrality of the molecule;  $\text{R}_{2\text{a-c}}$   
 and  $\text{R}_{3\text{a-d}}$  are hydrogen,  $\text{R}_1$  is methyl; X is oxygen, Z is  
 10 omitted, and Y is a polysubstituted aryl moiety of Formula  
 III where  $\text{R}_4$  and  $\text{R}_8$  are methyl,  $\text{R}_5$  and  $\text{R}_7$  are hydrogen,  
 and  $\text{R}_6$  is carboxyl ( $-\text{CO}_2\text{H}$ ). One of the more preferred  
 chemiluminescent substrate capable of reverse emission has  
 15 structure **13**. It is readily converted by alkaline  
 phosphatase to its keto form (**14**).

**13****14**

Another preferred chemiluminescent substrate capable of reverse emission is the reduced form of compound **13**,

**15****16**

which is shown as **15**. Similar to **13**, compound **15** is also readily converted by alkaline phosphatase to its keto form (**16**).

It has been unexpectedly discovered that both compounds **13** and **15** are capable of emitting light at extremely long wavelength. Figure 2M is the emission spectrum of compound **15** determined by FFS, showing that it emits light at  $\lambda_{\text{max}}$  678 nm. Figure 2N is the emission spectrum of compound **16**, which is the product of **15** after the treatment with AP. It shows that **16** emits light at  $\lambda_{\text{max}}$  450 nm, giving a net 228 nm of hypsochromic shift from **15**.

Hypsochromic shift of emission wavelength due to a number of factors including solvent and chemical substitutional effects on the acridinium compounds has been disclosed in the co-pending WO 00/09487, which is included herein as reference. Distinguishably, one of the discoveries in this invention which relates to a large


-51-

hypsochromic shift of emission wavelength from 13 to 14 and from 15 to 16 is due to the interruption of electronic conjugation between the acridinium nucleus and the electron-rich aromatic side chain caused by the enzymatic  
5 dephosphorylation.

## 2. Substrates and products having different emission kinetics

In addition to spectral distinction as a means of  
10 discriminating chemiluminescent substrate from product, enzyme-mediated, alteration of acridinium-ester flash-kinetics is also an effective method to monitor and determine the concentration of the enzyme. This methodology thus provides an alternate 'readout' signal  
15 when the enzyme is used as the label in an assay. One approach to modulate acridinium ester light-emission kinetics is to alter the electronic properties of a suitable functional group located on the phenol, as shown in Reaction E. During the chemiluminescent reaction of  
20 acridinium esters, cleavage of the phenolic ester is a prerequisite for formation of the dioxetanone precursor that is ultimately responsible for light emission. If the cleavage of this phenolic ester can be altered in a predictable way by making the phenol a better or poorer  
25 leaving group, then the kinetics of light emission from the corresponding acridinium ester can be modulated. For instance, an electron-donating group located ortho or para to the phenolic hydroxy moiety will render the phenol electron-rich hence a poor leaving group. This will lead  
30 to slow light emission. Conversion of the electron-

donating group to an electron-withdrawing group will accelerate cleavage of the phenolic ester and fast light emission will be obtained. The conversion of an electron-donating to an electron-withdrawing group or vice versa can be accomplished enzymatically. For instance, it is well known that oxidative enzymes such as peroxidases can oxidize electron-donating group such as amino to electron-withdrawing groups such as nitroso or nitro groups. On the other hand, hydrolytic enzymes can convert an electron-withdrawing or neutral functional group to an electron-donating group. In this case the kinetics of light emission from an acridinium ester is slowed down by the enzyme label.



Slow light emission

Fast light emission

EDG = electron-donating group


EWG = electron-withdrawing group

15

Reaction E


Acridinium esters containing a 4'-hydroxy group on the phenol exhibit slow kinetics in their light emission upon reaction with hydrogen peroxide in strong alkaline solution. This is primarily because the phenol in the acridinium ester containing the 4'-hydroxy group is a poor

leaving group. Conversion of this hydroxy group to a phosphate ester attenuates somewhat the electron donating ability of the 4'-oxygen substituent. Light emission from acridinium esters containing a 4'-phospho substituent is relatively faster than their 4'-hydroxy counterparts. Within this category are a subgroup of chemiluminescent acridinium substrates that are capable of emitting light at slower kinetics after the treatment with a hydrolytic enzyme, said subgroup of chemiluminescent acridinium substrates having Formula XIII. Wherein,  $R_1$ ,  $R_{2a-c}$ ,  $R_{3a-d}$ ,  $A^-$ ,  $M$ , and  $P$  are as defined in Formula II,  $R_{2d}$  is as defined for  $R_{2a}$ ,  $R_{2b}$ ,  $R_{3b}$ , and  $R_{3d}$ , and  $R_5$  and  $R_7$  are defined in Formula III.  $R_{16}$  and  $R_{17}$ , identical or different, are the groups selected from hydrogen, methyl, alkyl with low molecular weight, and halides. Preferably,  $R_{16}$  and  $R_{17}$  are different and one of them is hydrogen. More preferably, both  $R_{16}$  and  $R_{17}$  are hydrogen.



Formula XIII

One of the more preferred chemiluminescent acridinium substrates of Formula XIII is compound 4'-Phos-AE (17). As shown in Reaction F, 17 is readily converted by AP to the product 4'-OH-AE (18). It was found that at short measuring times (0.5-0.3 s), 17 emitted light ~190 times faster than 18. While theoretically, either the concentration of the substrate or the product can be measured to estimate enzyme activity, it was found to be more convenient (for effective signal discrimination) to measure the chemiluminescent activity of the substrate, 4'-Phos-AE (17). The chemiluminescent response to the concentration of alkaline phosphatase using 17 as the substrate is given in Figure 13.



Reaction F

### 3. Light Emission Spectra

The light emission spectra of compounds 1 ~ 12 and 15~18 were determined by a Fast Spectral Scanning System (FSSS) of Photo Research (a division of Kollmorgen Corp) of Burbank, Calif., U.S.A. The experiment was carried out in a

dark room. Each compound was dissolved in acetonitrile or *N,N*-dimethylformamide. The resulting concentrate was diluted with the same solvent to form the working solution, which upon flashing gave a light emission with an adequate intensity. A typical experiment utilized 10 ~ 100 µg or more of the sample in 500 µl of the solvent contained in a 13 x 100 mm borosilicate test tube. The tube was placed on a test tube rack raised to a proper height. A piece of aluminum foil was placed on the back of the tube to enhance the detectability of the emitted light. The FSSS optical head was placed in front of the tube at an approximate distance of about 130 mm with its lens focused on the liquid in the tube. The sample solution was first treated with 0.35 ml of the Flashing Reagent #1 (Bayer Diagnostics) containing 0.1 N HNO<sub>3</sub> and 0.5% H<sub>2</sub>O<sub>2</sub>. The room was then darkened, and 0.35 ml of the Flashing Reagent #2 (Bayer Diagnostics) containing 0.25 N NaOH and 0.5% ARQUAD was added to the reaction mixture immediately. (See U.S. Pat. No. 4,927,769 which is commonly assigned and incorporated herein by reference.) The light which was generated instantaneously following the addition of the Reagent #2 was recorded by FSSS for 5 seconds starting from about one second before the Reagent #2 was added. The various emission spectra determined on FSSS are given in Figures 2A-2P, and also summarized in Table 1.



Table 1

	Compound	Emission	
		Range* (nm)	Maximum (nm)
5	1	430—600	478
	2	500—780^	602
	3	430—600	474
	4	500—780^	604
10	5	430—600	478
	6	500—780^	600
	7	430—600	474
	8	500—780^	594
	9	440—600	476
15	10	520—780^	674
	11	430—600	474
	12	510—780^	638
	15	520—780^	678
	16	410—580	450
20	17	410—540	436
	18	410—540	452

\* Range is set for spectral region with signal intensity of above 5% of peak height.

^ Emission spectral range goes beyond the scanning limit (380-780 nm) of FSSS.

#### 4. Applications of the chemiluminescent enzyme substrates in binding assays

While the specific example of the actual diagnostic assay disclosed here uses 2-Phos-DMAE (1) as a chemiluminescent substrate for alkaline phosphatase, where alkaline phosphatase serves as a label for the detection of human thyroid stimulating hormone (TSH) in serum, it is reasonable to conclude that a variety of hydrolytic enzymes could be used in conjunction with the appropriate chemiluminescent substrates in a variety of assay architectures for detection of either endogenous, diagnostic enzyme markers or other clinically relevant diagnostic markers should the hydrolase be used as a label. Therefore, while we claim those assay architectures as should be obvious or otherwise apparent to those who are skilled in the art of enzymatic, diagnostic assay, we do not restrict our claims to the following descriptions.

a. Conversion of assay readout systems from colorimetric or fluorescence detection to chemiluminescence detection:

The co-application of enzyme labels and their dependent luminescent chemistries to bioanalytical techniques has been reported as one of several strategies for developing ultrasensitive clinical detection methodologies [Clin. Biochem., 26, 325, (1993)]. Conversion of existing diagnostic test formats from their current methods of colorimetric or fluorometric measurement to the potentially more sensitive method of chemiluminescence detection would

have a diverse multiplicity of application in clinical analysis, especially since these colorimetric or fluorometric assays already exist in multitudinous configurations. As mentioned earlier, 5 chromogenic and fluorometric indicator substances have been used in sensitive immunoassays in the form of enzyme substrates for detection of enzymes or the analytes to which these enzymes are directed as labels. The present invention discloses an analytical procedure where in general the 10 chemiluminescent-emission properties of an enzyme substrate are distinguishable from the resultant product and that the enzymatic source for the catalytic conversion of said substrate to product can be used to measure the enzyme activity directly for 15 measurement of either the enzyme or the analytes to which the enzyme would be attached as a label.

b. An analyte, as in the example below on human TSH, is complexed specifically with two different 20 antibodies immobilized on solid phase and tagged with an enzyme label, respectively. The quantity of analyte in a sample should correlate to the quantity of captured label, which in this case is alkaline phosphatase, and should ultimately correlate to the 25 observed magnitude of chemiluminescence. However, we envision more broadly the following scheme of assay architectures generally divided into the two classes referred to independently as immunoassays and nucleic acid assays. We further divide these classes into 30 either homogeneous or heterogeneous assay

configurations dependent on the separation of bound from free analyte. Furthermore we do not restrict our claims to those assay systems currently employing enzymes as labels, i.e. EIA, ELISA, Emit®, etc., but encompass also the realm of clinical diagnostics for which enzymes may be freely substituted for non-enzyme labels, i.e. radioisotopes, chromaphores, fluores, etc.

c. Heterogeneous Chemiluminescent Assay

An enzyme captured by a solid phase matrix, for example, a polystyrene surface, magnetic particle, or combination of both, or precipitating protein, etc., either non-specifically, as with adsorption, or specifically as through the attachment to the solid phase, non-covalently or otherwise, of a [directed] binding partner, including complementary nucleic acid sequence, biotin, antibody, binding protein, receptor, ligand, etc. would be separated from interfering substances through the separation of the solid phase from other assay components prior to application of the chemiluminescent substrate.

The enzyme could be either the endogenous diagnostic marker of interest, a label attached, covalently or otherwise, to a specific binding partner (collectively termed a tracer or probe) which is captured by the solid phase or a secondary reagent required for signal generation or amplification as in enzyme-cycling assays.

-60-

Examples of these assay configurations are widely reported in the literature and compendiums of specific constructs have been published (Maggio, E.

Enzyme-immunoassay; (1990) CRC Press) (Wild, D.;  
5 The Immunoassay Handbook; (1994); Stockton Press).

A heterogeneous enzyme-immunoassay for the detection of an enzyme phosphatase might be set up in several configurations. The simplest method would be to capture the phosphatase using an antibody which  
10 would specifically bind the phosphatase to separable solid phase, washing interfering substances from the solid phase prior to the addition of chemiluminescent phosphorylated substrate. A second method directed toward a protein analyte, could be fashioned in which  
15 a sandwich assay utilizing two antibodies, the first of which would be linked to a solid phase as described above, the second would be coupled to the signal generating system, alkaline phosphatase,  $\beta$ -galactosidase etc. The concentration of the enzyme  
20 label in the assay is then measured by application of the specific chemiluminescent substrate. Alternatively, in a competitive assay for a small analyte, a chemiluminescence generating enzyme such as alkaline phosphatase can be coupled to the same  
25 hapten for use in the assay as a tracer, where endogenous analyte and the hapten-enzyme conjugate would compete for a limited number of solid phase binding sites using a similar signal readout mechanism as is described above for the sandwich  
30 assay.

-61-

d. Homogeneous chemiluminescent assays

Assays that employ a homogeneous format can also be coupled to the present invention. Homogeneous assay architectures do not require the separation of assay components to discriminate between negative and positive analyte controls and therefore require fewer processing steps than heterogeneous assays. Homogeneous assay technologies such as Emit® and CEDIATM could be adapted to chemiluminescence detection. Emit® assays for small analytes could be developed in which both a hapten-enzyme conjugate, similar to that described above, and an analyte specific antibody would be added to a clinical sample. The resultant chemiluminescence would be dependent on the degree of enzyme inhibition occurring with the formation of the antibody-enzyme complex. CEDIATM assays are envisioned in which a hapten is covalently conjugated to the amino-terminal fragment of recombinant  $\beta$ -galactosidase or the enzyme donor (ED) fragment. Both the hapten-ED conjugate and an analyte specific antibody would then be incubated with the clinical sample in the presence of the carboxy-terminal fragment of  $\beta$ -galactosidase, termed enzyme acceptor (EA) (Engel, W.; Khanna, P.; J. Immunol. Methods: (1992); 150; 99). The magnitude of resultant chemiluminescence from the supposed enzyme substrate catalysis product, 2-hydroxy-DMAE, would depend on the degree of remaining enzyme inactivation.

-62-

## e. Chemiluminescent nucleic acid assays


We propose as obvious the application of said chemiluminescent enzyme substrate systems for the labeling and /or detection of nucleic acids in nucleic acid assays.

The invention disclosed herein is illustrated, but not limited, by the following examples.

Example 1.

Synthesis of (2',6'-dimethyl-4'-benzyl-oxycarbonyl)phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate trifluoroacetate (2-OH-DMAE-Bn, 4).

-63-



#### 4-Methoxyethoxymethoxy-iodobenzene

- 5 A solution of 4-iodophenol (10 g, 45.45 mmol) in 200 ml of anhydrous tetrahydrofuran was treated at 0°C with sodium hydride (2.36 g, 60% dispersion, 59.09 mmol) for 5 minutes. To the resulting mixture was slowly added over a 5-minute period methoxyethoxymethyl chloride (8.3 ml,



-64-

72.73 mmol). The mixture was stirred at 0°C under nitrogen for 30 minutes, warmed to room temperature, and stirred for 24 hours. The solvent was removed under reduced pressure. The residue was taken into 500 ml of ether, washed with 5% sodium hydroxide (4 x 200 ml), water (4 x 200 ml), saturated sodium chloride (1 x 200 ml), and dried over sodium sulfate. Evaporation of the solvent under reduced pressure gave an oily product in 14.1 g. TLC (silica gel, ether): Rf 0.5.

10

N-(4-Methoxyethoxymethoxy)phenyl isatin

A solution of isatin (4.0 g, 27.2 mmol) in 200 ml of anhydrous N, N-dimethylformaldehyde was treated at room temperature with sodium hydride (1.036 g, 60% dispersion, 32.64 mmol) for 0.5 hour, followed by addition of 4-methoxyethoxymethoxy-iodobenzene (12.57 g, 40.8 mmol) and copper (I) iodide (10.34 g, 54.4 mmol). The resulting mixture was stirred at 160°C under nitrogen for 17 hours. It was cooled to room temperature, and diluted with 400 ml of chloroform. The resulting mixture was filtrated to remove the inorganic materials. The filtrate was evaporated under reduced pressure to give a crude mixture containing N-(4-methoxyethoxymethoxy)phenyl isatin as a major product. TLC (silica gel, ether): Rf 0.8.

25

2-Methoxyethoxymethoxy-acridine-9-carboxylic acid

The above crude 4-methoxyethoxymethoxyphenyl isatin, without purification, was suspended in 120 ml of 10% potassium hydroxide. The suspension was refluxed at 150°C for 5 hours. After cooling to room temperature, the

30

-65-

mixture was filtrated to remove the orange impurities. The filtrate was acidified in an ice-water bath with concentrated hydrochloric acid to pH 2. The resulting yellow precipitate was collected and washed with water (4  
5 x 50 ml) and air-dried. The dried material was further washed with ether (6 x 50 ml) to yield the desired product in 6.7 g. TLC (silica gel, 30% methanol/chloroform): Rf 0.5.

10 (2',6'-Dimethyl-4'-benzyloxycarbonyl)phenyl 2-methoxy-ethoxy-methoxy-acridine-9-carboxylate

A suspension of 2-methoxyethoxymethoxy-acridine-9-carboxylic acid (3.6 g, 11 mmol) in 150 ml of anhydrous pyridine was treated with p-toluenesulfonyl chloride  
15 (4.183 g, 22 mmol) at 0°C for 5 minutes to form a homogeneous brown solution. Then, benzyl 3,5-dimethyl-4-hydroxy-benzoate (2.818 g, 11 mmol) was added. The solution was stirred at room temperature under nitrogen for 20 hours. The solvent was removed under reduced  
20 pressure. The residue was separated on a silica flash chromatography column packed in hexane. It was eluted with 50% ether/hexane (1 liter) followed by 70% ether/hexane (3 liters). The product fraction was obtained from the 70% ether/hexane eluent. Evaporation of  
25 the solvents under reduced pressure gave 3.74 g of the desired product. TLC (silica gel, ether): Rf 0.8.


(2',6'-Dimethyl-4'-benzyloxycarbonyl)phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate trifluoroacetate

A light-yellow solution of (2',6'-dimethyl-4'-benzyloxycarbonyl)phenyl 2-methoxyethoxymethoxy-acridine-9-carboxylate (400 mg, 0.708 mmol) in 20 ml of anhydrous methylene chloride was treated with methyl trifluoromethanesulfonate (0.4 ml, 3.54 mmol) at room temperature under nitrogen with stirring for 14 hours. The resulting mixture was treated with anhydrous ether (20 ml). The precipitate was collected and washed with ether (4 x 20 ml) to yield 325 mg of the crude product. MS: (ESI): m/z 492.6 (M<sup>+</sup>). <sup>1</sup>H NMR (300 MHz, MeOD-d<sub>4</sub>/CDCl<sub>3</sub>): δ 2.52 (6H, s), 5.01 (3H, s), 5.42 (2H, s), 7.37 - 7.51 (5H, m), 7.81 (1H, d, J = 2.6 Hz), 7.97 (2H, s), 8.10 (1H, t, J = 7.0 Hz), 8.16 (1H, dd, J<sub>1</sub> = 8.0 Hz, J<sub>2</sub> = 2.6 Hz), 8.42 (1H, t, J = 7.0 Hz), 8.60 (1H, d, J = 8.8 Hz), and 8.73 (2H, two overlapping doublets, J ~ 8.0 Hz). The product (25 mg) was further purified on a preparative HPLC column (YMC, 250 x 30 mm, ODS, 10 μm), eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TAF/acetonitrile (solvent B) in the following manner: 40 to 60% B in 40 minutes, flow rate 20 ml/minute, monitored at 260 nm. The desired product at retention time of ~30 minutes was collected and crystallized from methylene chloride/ether to give 17 mg of pure 4.

**Example 2.**

Synthesis of (2',6'-dimethyl-4'-benzyloxy-carbonyl)phenyl 2-phosphoryloxy-10-methyl-acridinium-9-carboxylate trifluoroacetate (2-Phos-DMAE-Bn, 3)

5



A<sup>-</sup> = CF<sub>3</sub>CO<sub>2</sub><sup>-</sup> after the compound was recovered from HPLC mobile phase containing CF<sub>3</sub>CO<sub>2</sub>H

A solution of (2',6'-dimethyl-4'-benzyloxycarbonyl)phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate trifluoromethane-sulfonate (50 mg, 0.102 mmol) in pyridine (0.5 ml) was cooled to 0°C and then added dropwise over a 3 minute period to a pre-cooled solution of phosphorous oxychloride (57 µl, 6 eq.) in 0.5 ml of pyridine. The reaction mixture was stirred at 0°C under nitrogen for 30 minutes. The reaction was quenched with 400 µl of 0.5 N NaOH followed by another 200 µl of 1 N NaOH. The mixture was diluted with 1 ml of water, and then filtrated. The resulting yellow solid was dissolved in mixed DMF and water, and separated on a preparative HPLC column (YMC, 250 x 30 mm, ODS, 10 µm), eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TAF/acetonitrile (solvent B) in the following manner: 30 to 60% B in 40


minutes, flow rate 20 ml/minute, monitored at 260 nm. The desired product (3) at retention time of 32 minutes was obtained in 20 mg. MS (MALDI-TOF): m/z 573 (M + 1).

5

**Example 3.**

**Synthesis of (2',6'-dimethyl-4'-carboxyl)phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate trifluoroacetate (2-OH-DMAE, 2)**

10




A<sup>-</sup> = CF<sub>3</sub>CO<sub>2</sub><sup>-</sup> after the compound was recovered from HPLC mobile phase containing CF<sub>3</sub>CO<sub>2</sub>H

A solution of (2',6'-dimethyl-4'-benzyloxycarbonyl)phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate trifluoromethanesulfonate (100 mg) in 4 ml of 30% hydrogen bromide in acetic acid was stirred at 55°C under nitrogen for 1 hour, and then treated with 10 ml of anhydrous ether. The resulting precipitate was collected and washed with ether (4 x 10 ml) to give 80 mg of (4-carboxyl-2,6-dimethyl)phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate bromide. MS (ESI): m/z 402.7 (M<sup>+</sup>). <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>CN/MeOD-d<sub>4</sub>): δ 2.52 (6H, s), 4.95

(3H, s), 7.78 (1H, d,  $J = 2.7$  Hz), 7.76 (2H, s), 8.10 (1H, t,  $J = 7.0$  Hz), 8.13 (1H, dd,  $J_1 = 9.9$  Hz,  $J_{3=2} = 2.7$  Hz), 8.40 (1H, d t,  $J_1 = 2.7$  Hz,  $J_2 = 8.0$  Hz), 8.62 (1H, d,  $J = 8.0$  Hz), 8.77 (1H, d,  $J = 9.2$  Hz), and 8.78 (1H, d,  $J = 9.9$  Hz). The further purification of the above product (68 mg) was carried out by preparative HPLC (YMC, 250 x 30 mm, ODS, 10  $\mu$ m), eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TAF/acetonitrile (solvent B) in the following manner: 10 to 60% B in 40 minutes, flow rate 20 ml/minute, monitored at 260 nm. The desired product (2) at retention time of ~27 minutes was obtained in 46 mg.

#### 15 Example 4.

Synthesis of (2',6'-dimethyl-4'-carboxyl)phenyl 10-methyl-2-Phosphoryloxy-acridinium-9-carboxylate bromide (2-Phos-DMAE, 1)




20

A mixture of (2',6'-dimethyl-4'-benzyloxycarbonyl)phenyl 10-methyl-2-phosphoryloxy-10-methyl-acridinium-9-carboxylate trifluoroacetate (3, 37 mg) and 2.5 ml of 30% hydrogen bromide in acetic acid was

stirred at 50°C under nitrogen for 1.5 hour. The resulting mixture was cooled to room temperature, and treated with ether to form yellow precipitate. It was dissolved in mixed DMF/water with the help of a small amount of triethylamine. The mixture was separated on a preparative HPLC column (YMC, 250 x 30 mm, ODS, 10  $\mu$ m), eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TAF/acetonitrile (solvent B) in the following manner: 10 to 60% B in 40 minutes, flow rate 20 ml/minute, monitored at 260 nm. The fraction at retention time of ~17 minutes was collected and lyophilized to dryness to give pure 1 in 5.3 mg. MS (DALTI-TOF): m/z 482.818 (M + 1).

15 **Example 5.**

**Synthesis of phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate trifluoromethanesulfonate (2-OH-AE, 6)**



Phenyl 2-methoxyethoxymethoxy-acridine-9-carboxylate

To a solution of 2-methoxyethoxymethoxy-acridine-9-carboxylic acid (101 mg, 0.308 mmol) in 5 ml of anhydrous pyridine was added *p*-toluenesulfonyl chloride (117 mg, 0.616 mmol). It was stirred at 0°C for 5 minutes and at room temperature for additional 10 minutes before 29 mg (0.038 mmol) of phenol was added. The reaction was stirred at room temperature under nitrogen overnight. The reaction mixture was evaporated under reduced pressure. The residue was suspended in mixed methanol and ethyl acetate and filtrated. The resulting filtrate was reduced to a small volume and separated on 4 preparative silica gel plates (20 x 20 cm x 2 mm thick), which was developed with hexane/ether (2 : 1). The major product band was collected and eluted with the same solvent system. Removal of the solvents gave 27 mg of the pure product. TLC (silica gel, hexane/ether 2 : 1): R<sub>f</sub> 0.9.


Phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate tri-fluoromethanesulfonate

A solution of phenyl 2-methoxyethoxymethoxy-acridine-9-carboxylate (25 mg, 0.062 mmol) in 1.5 ml of anhydrous methylene chloride was treated with methyl trifluoromethanesulfonate (38 µl, 0.336 mmol) at room temperature under nitrogen with stirring overnight. The reaction was diluted with another 0.5 ml of methylene chloride and then treated with anhydrous ether (4 ml). The resulting precipitate was collected and washed with ether to give 6 in 19 mg. MS (ESI): m/z 330 (M<sup>+</sup>).



**Example 6.**

**Synthesis of phenyl 10-methyl-2-Phosphoryloxy-acridinium-9-carboxylate trifluoroacetate (2-Phos-AE, 5)**



A<sup>-</sup> = CF<sub>3</sub>CO<sub>2</sub><sup>-</sup> after the compound was recovered from HPLC mobile phase containing CF<sub>3</sub>CO<sub>2</sub>H


5

A solution of phenyl 2-hydroxy-10-methyl-acridinium-9-carboxylate trifluoro-methanesulfonate (19 mg, 0.040 mmol) in pyridine (0.3 ml) was cooled to 0°C and then added slowly to a pre-cooled solution of phosphorous oxychloride (22 µl, 0.24 mmol) in 0.3 ml of pyridine. The reaction mixture was stirred at 0°C under nitrogen for 30 minutes, and quenched with 500 µl of 5% ammonium hydroxide for 10 minutes. The solution was then diluted with 1 ml of water and neutralized with 1N HCl. The mixture was separated on a preparative HPLC column (YMC, 300 x 20 mm, ODS, 10 µm), eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TAF/acetonitrile (solvent B) in the following manner: 10 to 60% B in 40 minutes, flow rate 20 ml/minute, monitored at 260 nm. The desired product (5) at retention time of 18 minutes was obtained in 8 mg. MS (ESI): m/z 410 (M<sup>+</sup>).

**Example 7.**

**Synthesis of (2', 6'-dimethyl-4'-carboxyl)phenyl 2-phosphoryloxy-10-sulfobutyl-acridinium-9-carboxylate (2-Phos-NSB-DMAE, 7)**

5



(2', 6'-Dimethyl-4'-benzyloxycarbonyl)phenyl 2-hydroxyacridine-9-carboxylate

10 A solution of (2', 6'-dimethyl-4'-benzyloxy-carbonyl)phenyl 2-methoxyethoxymethoxy-acridine-9-carboxylate (960 mg, 1.7 mmol) in methylene chloride (5 ml) was treated with trifluoroacetic acid (5 ml) at room temperature for 19 hours. The reaction mixture was blown to

dryness with a nitrogen stream and the residue was suspended in ether. The product was collected and further washed with ether (2 x 10 ml) to give 610 mg. MS (MALTI-TOF): m/z 479 (M + 1).

5

(2', 6'-Dimethyl-4'-benzyloxycarbonyl)phenyl 2-dimethyl-phosphoryloxy-acridine-9-carboxylate

To a solution of (2', 6'-dimethyl-4'-benzyloxycarbonyl)phenyl 2-hydroxy-acridine-9-carboxylate  
10 (50 mg, 0.105 mmol) in pyridine (2 ml) at 0°C under nitrogen was added sodium hydride (5 mg, 0.208 mmol). It was allowed to stir at room temperature for 1 hour before dimethoxychlorophosphate (56 µl, 0.519 mmol) was added. The reaction was continued to stir for 3 hours. The  
15 product was isolated from preparative HPLC column (YMC, 250 x 30 mm, ODS, 10 µm). The column was eluted in gradient by mixing 0.05% TFA/water (solvent A) and 0.05% TFA/acetonitrile (solvent B) in the following manner: 40 to 80% B in 40 minutes, flow rate at 20 ml/minute,  
20 monitored at 260 nm. The fraction at retention time of ~52 minutes was collected and lyophilized to dryness to give the desired product in 45 mg. MS (MALTI-TOF): m/z 587 (M + 1).

25 (2', 6'-Dimethyl-4'-benzyloxycarbonyl)phenyl 2-dimethyl-phosphoryloxy-10-sulfobutyl-acridinium-9-carboxylate

A mixture of (2', 6'-dimethyl-4'-benzyloxycarbonyl)phenyl 2-dimethylphosphoryloxy-acridine-9-carboxylate (45 mg, 0.0768 mmol) and 1,4-butanedisulfone (1

ml, 9.78 mmol) was stirred at 150°C for 6 hours under nitrogen. The resulting thick gum was dissolved in a mixture of acetonitrile/water. The solution was separated on a preparative HPLC column (YMC, 250 x 30 mm, ODS, 10  
5  $\mu$ m). The column was eluted in gradient by mixing 0.05% TFA/water (solvent A) and 0.05% TFA/acetonitrile (solvent B) in the following manner: 40 to 80% B in 40 minutes, flow rate at 20 ml/minute, monitored at 260 nm. The fraction at retention time of ~21 minutes' was collected  
10 and lyophilized to dryness to give the desired product in 1 mg. MS (MALTI-TOF): m/z 723 (M + 1).


(2', 6'-Dimethyl-4'-carboxyl)phenyl 2-phosphoryloxy-10-sulfo-butyl-acridinium-9-carboxylate

15 A solution of (2', 6'-dimethyl-4'-benzyl-oxycarbonyl)phenyl 2-dimethylphosphoryloxy-10-sulfo-butyl-acridinium-9-carboxylate (1 mg, 0.00140 mmol) in chloroform (0.5 ml) was treated with boron tribromide (4  $\mu$ l, 0.0423 mmol). It was stirred at room temperature under  
20 nitrogen for 3 hours. The mixture was blown with a nitrogen stream to dryness then dissolved in acetonitrile and water mixture. The desired product was isolated from a semi-preparative HPLC column (Phenomenex, 300 x 7.8 mm, ODS, 10  $\mu$ m). The column was eluted in gradient by mixing  
25 0.05% TFA/water (solvent A) and 0.05% TFA/acetonitrile (solvent B) in the following manner: 10 to 60% B in 40 minutes, flow rate at 2.5 ml/minute, monitored at 260 nm. The fraction at retention time of ~17 minutes was

collected and lyophilized to dryness to give the desired product in 0.1 mg. MS (MALTI-TOF):  $m/z$  605 ( $M + 1$ ).

5 **Example 8.**

Synthesis of (2', 6'-dimethyl-4'-carboxyl)phenyl 2-hydroxy-10-sulfoethyl-acridinium-9-carboxylate (2-OH-NSB-DMAE, 8)



-77-

N-(4-methoxyphenyl)isatin

A solution of isatin (2.5 g, 0.017 mol) in anhydrous DMF (50 ml) was cooled to 0°C under a nitrogen atmosphere and treated with sodium hydride (0.5 g, 1.2 equivalents). A purple solution was formed which was stirred at 0°C for 30 minutes and then warmed to room temperature. 4-Bromoanisole (2.13 ml, 1 equivalent) was added followed by copper iodide (6.46 g, 2 equivalents). The reaction was heated in an oil-bath at 145 °C for 7 hours. The reaction was then cooled to room temperature and diluted with an equal volume of ethyl acetate. This suspension was filtered and the filtrate was evaporated to dryness. TLC (1:4, ethyl acetate:hexanes) indicated a very clean reaction; R<sub>f</sub> (product) = 0.5. The crude material was used as such for the next reaction.

2-Methoxy-acridine-9-carboxylic acid

The crude N-(4-methoxyphenyl)isatin from the above was suspended in 10% aqueous potassium hydroxide (150 ml) and refluxed under a nitrogen atmosphere. After 4 hours, the reflux was stopped and the reaction was filtered while still hot. The filtrate was diluted with water (~150 ml) and ice. This solution was then acidified with concentrated hydrochloric acid. A yellow precipitate appeared which was collected by filtration. The precipitate was rinsed with cold water and ether and then air-dried. The dried residue was then transferred to a round-bottom flask with the aid of methanol and the partial solution was evaporated to dryness. The resulting residue was evaporated to dryness from toluene twice. A

yellowish-brown powder was recovered in 1.15 g. TLC (1:4, methanol:chloroform) indicated a clean reaction; R<sub>f</sub> (product) = 0.14. This material was used as such for the next reaction.

5

2',6'-Dimethyl-4'-benzyloxycarbonylphenyl 2-methoxyacridine-9-carboxylate

2-Methoxyacridine-9-carboxylic acid (0.8 g, 0.0032 mol) in anhydrous pyridine (50 ml) was cooled in an ice-bath under a nitrogen atmosphere and treated with *p*-toluenesulfonyl chloride and 4-benzyloxycarbonyl-2,6-dimethylphenol (0.81 g, 0.0032 mol). The reaction was warmed to room temperature and stirred under a nitrogen atmosphere for 24 hours. The solvent was then removed under reduced pressure and the residue was dissolved in chloroform (10 ml). The product was purified by flash chromatography using 5% ethyl acetate, 25% chloroform, 70% hexanes. Evaporation of the flash fractions containing the product yielded a bright yellow solid in 0.84 g. MS (MALDI-TOF): m/z MS 492.8 (M + 1).

(2', 6'-Dimethyl-4'-benzyloxycarbonyl)phenyl 2-methoxy-10-sulfobutyl-acridinium-9-carboxylate

A mixture of (2',6'-dimethyl-4'-benzyloxycarbonyl)phenyl 2-methoxy-acridine-9-carboxylate (200 mg, 0.407 mmol) in 2 ml of 1,4-butanedisulfone was stirred at 150°C for 19 hours under nitrogen. The resulting thick gum was dissolved in a mixed acetonitrile/water solvent. The solution was separated on a preparative HPLC column (YMC, 250 x 30 mm, ODS, 10 μm). The column was eluted in

-79-

gradient by mixing 0.05% TFA/water (solvent A) and 0.05% TFA/acetonitrile (solvent B) in the following manner: 30 to 60% B in 40 minutes, flow rate at 20 ml/minute, monitored at 260 nm. The fraction at retention time of  
5 ~37 minutes was collected and lyophilized to dryness to give the desired product in 58.5 mg. MS (MALTI-TOF): m/z 629 (M + 1).

10 (2', 6'-Dimethyl-4'-carboxyl)phenyl 2-hydroxy-10-sulfobutyl-acridinium-9-carboxylate

A solution of (2', 6'-dimethyl-4'-benzyl-oxy-carbonyl)phenyl 2-methoxy-10-sulfobutyl-acridinium-9-carboxylate (50 mg, 0.080 mmol) in chloroform (4 ml) was treated with boron tribromide (50 µl, 0.529 mmol) and was  
15 stirred at room temperature under nitrogen for 3 hours. The reaction was blown with a nitrogen stream to dryness then dissolved in acetonitrile and water mixture. The desired product was isolated from a preparative HLPC column (YMC, 250 x 30 mm I.D., ODS, 10 µm). The column was  
20 eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TFA/acetonitrile (solvent B) in the following manner: 10 to 60% B in 40 minutes, flow rate at 20 ml/minute, monitored at 260 nm. The fraction at retention time of ~26 minutes was collected and lyophilized to  
25 dryness to give the desired product in 18 mg. MS (MALTI-TOF): m/z 525 (M + 1). This compound was further purified, using another preparative HLPC column (YMC, 300 x 20 mm I.D., ODS, 10 µm). ). The column was eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05%  
30 TFA/acetonitrile (solvent B) in the following manner: 10




to 60% B in 40 minutes, flow rate at 16 ml/minute, monitored at 260 nm. The pure product was obtained in 4 mg. MS (MALTI-TOF):  $m/z$  525 ( $M + 1$ ).

5

**Example 9.**

Syntheses of (2',6'-dimethyl-4'-carboxyl)phenyl 2-hydroxy-7-methoxy-10-methyl acridinium-9-carboxylate (2-OH-7-MeO-DMAE, 10) and (2',6'-dimethyl-4'-carboxyl)phenyl 2-phosphoryloxy-7-methoxy-10-methyl acridinium-9-carboxylate (2-Phos-7-MeO-DMAE, 9)



Synthesis of 4-benzyloxybromobenzene

4-Bromophenol (2 g, 0.0116 mol) in acetone (40 ml) was treated with potassium carbonate (1.91 g, 1.2 equivalents) and benzyl bromide (1.44 ml, 1.05 equivalents). The reaction was refluxed under nitrogen. After 5-6 hours of reflux, the reaction was cooled to room temperature and diluted with an equal volume of ethyl acetate. This was diluted further with water and the organic layer was separated, dried over magnesium sulfate and evaporated to dryness to afford a white fluffy powder. Yield = 2.36 g (73%).

Synthesis of N-(4'-benzyloxy)phenyl-5-methoxyisatin

5-Methoxyisatin (1.5 g, 0.847 mmol) in anhydrous DMF (50 ml) was cooled in an ice-bath under nitrogen and treated with sodium hydride (0.25 g, 1.2 equivalents). After 15-20 minutes in ice, a solution of 4-benzyloxybromobenzene (2.36 g) in DMF (~3 ml) was added along with CuI (3.23 g, 2 equivalents). The resulting reaction was heated in an oil-bath at 140°C under nitrogen for 24 hours. The reaction was then filtered and the filtrate was evaporated to dryness. The residue was suspended in ethyl acetate and purified by flash chromatography using 35% ethyl acetate in hexane. Evaporation of the flash fractions afforded the alkylated istain as an orange-brown solid. Yield = 1 g (32%).

Synthesis of 2-benzyloxy-7-methoxyacridine-9-carboxylic acid

The N-alkylated isatin from above (1 g) was suspended in 10% potassium hydroxide (100 ml and refluxed under nitrogen for 4 hours. The reaction was then filtered while still hot and the filtrate was cooled in ice. This was acidified carefully with a mixture of ice and concentrated HCl till a thick yellow precipitate separated out. The precipitate was allowed to stand for ~15 minutes and was then collected by filtration. After rinsing with ether, the product was thoroughly air dried. Yield = 0.75 g (75%).

Synthesis of (2',6'-dimethyl-4'-benzyloxycarbonyl)phenyl 2-benzyloxy-7-methoxy-acridine-9-carboxylate

2-Benzyloxy-7-methoxy-acridine-9-carboxylic acid (0.36 g, 0.001 mol) in anhydrous pyridine (30 ml) was cooled in an ice-bath under nitrogen and treated with p-toluenesulfonyl chloride (0.39 g, 2 equivalents) followed by 4-carbobenzyloxy-2,6-dimethylphenol (0.3 g, 1.2 equivalents). The reaction was warmed to room temperature and stirred for 16 hours under nitrogen. The solvent was then removed under reduced pressure and the residue was dissolved in chloroform (~60 ml). The chloroform solution was washed with 3% aqueous ammonium chloride. It was then dried over magnesium sulfate and evaporated to dryness. The crude product was purified by preparative TLC using 70% hexane, 27% chloroform, 3% methanol and isolated as a yellow solid. Yield = 0.31 g (50%). MALDI-TOF MS 599.02 obs. (597.67 calc.).

Synthesis of (2',6'-dimethyl-4'-benzyloxycarbonyl)phenyl  
2-benzyloxy-7-methoxy-10-methyl-acridinium-9-carboxylate  
trifluoromethanesulfonate


The acridine ester from above (0.31 g, 0.52 mmol) was  
5 dissolved in dichloromethane (5 ml) and treated with  
methyl trifluoromethanesulfonate (0.575 ml, 10  
equivalents). The reaction was stirred at room temperature  
for 16 hours. Ether (150 ml) was then added and the  
precipitated product was collected by filtration and air  
10 dried. Yield = 0.23 g. MALDI-TOF MS 613.33 obs. (612.7  
calc.)

Synthesis of (2',6'-dimethyl-4'-carboxyl)phenyl 2-hydroxy-  
7-methoxy-10methyl-acridinium-9-carboxylate (10)

15 The acridinium ester from above (0.124 g) was stirred  
in a mixture of methyl sulfide and 30% HBr/AcOH (1:1, 4  
ml). After 4 hours, ether was added to precipitate the  
product which was collected by filtration and air dried.  
This was dissolved in methanol and analyzed by analytical  
20 HPLC using a 3.9 x 300 mm c18 column and a 30 minute  
gradient of 10-100% acetonitrile/water each containing  
0.05% TFA at a flow rate of 1 ml/min and UV detection at  
260 nm. The product was found to elute at 15 minutes  
while the starting material eluted at 24 minutes.  
25 Approximately, 60% of the crude material was purified by  
preparative HPLC and the HPLC fractions were lyophilized  
to dryness. Yield = 42 mg (80%). MALDI-TOF MS 432.87  
obs. (432.45 calc.)

Synthesis of (2',6'-dimethyl-4'-carboxyl)phenyl 2-phosphoryloxy-7-methoxy-10-methyl-acridinium-9-carboxylate (9)

Crude deblocked acridinium ester from above (80 mg) was dissolved in pyridine (25 ml) and treated with phosphorus oxychloride (3 x 75  $\mu$ l, ~15 equivalents) at 0°C under nitrogen. The reaction was stirred for 1 hour and then quenched with water (3 ml) and stirred for an additional hour at room temperature. The reaction was then concentrated to a small volume. HPLC analysis using the same conditions as above but with a 40-minute gradient of 10-60% acetonitrile/water (each with 0.05% TFA) showed product eluting at 20 minutes with starting material eluting at 24 minutes. The product was isolated by preparative HPLC and the HPLC fractions were lyophilized to dryness. Yield = 3 mg yellow powder. MALDI-TOF MS 513.00 obs. (513.26 calc.)



**Example 10.****Synthesis of 2-OH-Spiroacridan (12) and 2-Phos-Spiroacridan (11)**5 **N-(4'-Benzyloxy)phenylisatin**

To a solution of isatin (4 g, 27.2 mmol) in DMF (50 ml) under nitrogen at room temperature was added NaH (871 mg, 34.5 mmol). The reaction color changed from orange to purple. It was stirred at room temperature for 30 min  
10 before 4-benzyloxyphenyl bromide (10.21 g, 38.8 mmol) and CuI (10.34 g, 54.4 mmol) were added. It was refluxed at 160 °C in an oil bath for 20 hours under nitrogen. After cooling to room temperature, the reaction was poured into chloroform (400 ml), and filtrated. The filtrate was  
15 concentrated to dryness under reduced pressure to give the desired product as a brown gum. It was used in the next step without further purification.

**2-Benzyloxyacridine-9-carboxylic acid**

20 The above mixture in 10% KOH/H<sub>2</sub>O (220 ml) was refluxed at 130°C for 20 hours. The reaction was filtered while warm, and the filtrate was cooled to 0°C before it was acidified with concentrated HCl to pH 3. The yellow precipitate was filtered, and the filter cake was washed  
25 with water (4 x 200 ml). It was dried under reduced pressure at 50°C for 20 hours. The desired product was obtained in 1.8 g. It was confirmed by MS (MALTI-TOF): m/z 331 (M + 1).

(2'-Benzyloxy)phenyl 2-benzyloxyacridine-9-carboxylate

A solution of 2-benzyloxyacridine-9-carboxylic acid (306 mg, 0.93 mmol) in pyridine (30 ml) was treated with *p*-toluenesulfonyl chloride (276 mg, 1.45 mmol) at room temperature under nitrogen for 30 min before 2-(benzyloxy)phenol was added. The reaction was stirred at room temperature for 20 hours. It was concentrated to dryness under reduced pressure. The resulting material was purified on flash column eluted with gradient solvent system of ethyl acetate / hexane starting at 10%. The product came out at 20%. The desired fractions were combined and concentrated to dryness under reduced pressure to give 226 mg of the desired product. It was confirmed by MS (MALTI-TOF):  $m/z$  513 ( $M + 1$ ).

15

(2'-Benzyloxy)phenyl 2-benzyloxy-10-methyl-acridine-9-carboxylate trifluoromethanesulfonate

A solution of (2'-benzyloxy)phenyl 2-benzyloxyacridine-9-carboxylate (109 mg, 0.213 mmol) in dichloromethane (5 ml) was treated with methyl trifluoromethanesulfonate (260  $\mu$ L, 2.30 mmol) under nitrogen at room temperature for 20 hours. The reaction was blown to dryness with nitrogen followed by suspending in ether. The yellow precipitate was washed with more ether (4 x 10 ml). The resulting solid was dried under reduced pressure to give 71.35 mg of the desired product.

25

2-OH-Spiroacridan (11)

A mixture of (2'-benzyloxy)phenyl 2-benzyloxy-10-methyl-acridine-9-carboxylate (30 mg, 0.043 mmol) in 30%

30

HBr/AcOH (400  $\mu$ L) was stirred at 45 °C for 1 hour. It was blown with nitrogen to dryness followed by suspending in ether. The solid was filtered, washed with more ether (4 x 10 ml), and dried under reduced pressure to give 17 mg of the desired product. A portion of this product (11 mg) was purified on a preparative HLPC column (YMC, 250 x 20 mm I.D., ODS, 10  $\mu$ m). The column was eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TFA/acetonitrile (solvent B) in the following manner: 10 to 60% B in 40 minutes, flow rate at 16 ml/minute, monitored at 260 nm. The fraction at retention time of ~25 minutes was collected and lyophilized to dryness to give the pure product in 7.3 mg. It was confirmed by MS (MALTI-TOF): m/z 347 (M + 1).

15

#### 2-Phos-Spiroacridan (12)

To a solution of the above 2-OH-Spiroacridan (5.66 mg, 0.016 mmol) in pyridine (0.5 ml) at 0°C under nitrogen was added POCl<sub>3</sub> (7.65  $\mu$ L, 0.082 mmol). It was stirred at the same temperature for 1 hour then quenched with water (0.5 ml) and stirred for another 15 min. It was concentrated to dryness under reduced pressure, followed by purification on a preparative HLPC column (YMC, 250 x 20 mm I.D., ODS, 10  $\mu$ m). The column was eluted in gradient by mixing 0.05%TFA/water (solvent A) and 0.05% TFA/acetonitrile (solvent B) in the following manner: 10 to 60% B in 40 minutes, flow rate at 16 ml/minute, monitored at 260 nm. The fraction at retention time of ~18 minutes was collected and lyophilized to dryness to



give the pure product in 2.3 mg. It was confirmed by MS (MALTI-TOF):  $m/z$  427 ( $M + 1$ ).

5 Example 11.

Synthesis of (2',6'-dimethyl-4'-carboxyl)phenyl 3-( $\beta$ -phosphoryloxy-4'-hydroxystyryl)-10-methyl acridinium-9-carboxylate trifluoromethanesulfonate (3-Enol-Phos-DMAE, 13) and its corresponding acridan (3-Enol-Phos-acridan, 10 15)

